

GAUGED SUPERGRAVITIES FROM SPHERICAL REDUCTIONS

A Dissertation

by

TUAN ANH TRAN

Submitted to the Office of Graduate Studies of  
Texas A&M University  
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

December 2001

Major Subject: Physics

GAUGED SUPERGRAVITIES FROM SPHERICAL REDUCTIONS

A Dissertation

by

TUAN ANH TRAN

Submitted to Texas A&M University  
in partial fulfillment of the requirements  
for the degree of

DOCTOR OF PHILOSOPHY

Approved as to style and content by:

---

Christopher N. Pope  
(Chair of Committee)

---

Richard L. Arnowitt  
(Member)

---

Robert C. Webb  
(Member)

---

Stephen A. Fulling  
(Member)

---

Thomas W. Adair III  
(Head of Department)

December 2001

Major Subject: Physics

## ABSTRACT

Gauged Supergravities from Spherical Reductions. (December 2001)

Tuan Anh Tran, B.S., Hanoi University, Vietnam

Chair of Advisory Committee: Dr. Christopher N. Pope

This dissertation is devoted to deriving the bosonic sectors of certain gauged supergravities in various dimensions from reducing eleven-dimensional supergravity, type IIA and type IIB supergravities in ten dimensions on certain spherical spaces. Explicit non-linear Kaluza-Klein ansätze for reductions of eleven-dimensional supergravity and of type IIA and type IIB supergravities on  $S^n$  and  $S^n \times T^m$  are presented. Knowing explicit non-linear ansätze is proven to be very useful in finding super Yang-Mills operators of gauge theories via AdS/CFT correspondence. We present a sample calculation which allows us to find a super Yang-Mills operator using a non-linear ansatz. Knowing non-linear ansätze is also useful for finding supergravity duals to certain twisted supersymmetric gauge theories. These supergravity solutions are branes wrapped on certain supersymmetric cycles. Some solutions, which are dual to gauge theories in three and five dimensions, are presented.

To my parents for their love and their support

## ACKNOWLEDGMENTS

I am very grateful to my advisor, Christopher N. Pope, for not only teaching me Physics, but also for helping me with various issues in life. I am also grateful to Mirjam Cvetič and Hong Lü for the research we did together.

I am thankful to Glenn Agnolet, Richard Arnowitt, Michael Duff and Ergin Sezgin for teaching me various subjects of physics.

I am also thankful to Igor Lavrinenko, Mihail Mihailescu, Carlos Núñez, Inyong Park, Arta Sadrzadeh and Martin Schvellinger for the ideas they shared with me and for the work we did together.

Many stimulating discussions with Karim Benakli, Sadik Deger, Bashkar Dutta, Jianxin Lu, Sudipta Mukherji, Igor Rudychev, Hisham Sati, Per Sundell and Kaiwen Xu are acknowledged.

I would like to thank the people in the HEP section at ICTP, Trieste, Italy for teaching me HEP while I was in the Diploma Program. Particularly, I would like to thank Faheem Hussain and Seif Randjbar-Daemi for their help while I was at ICTP. Many conversations with K.S. Narain about aspects of string theory partially influenced my decision to work on supergravity and string theory. I would like to thank Goran Senjanović for his supervision while I was there.

I am indebted to Robert Foot and to Hoang Ngoc Long (my undergraduate advisor) for introducing me to particle physics.

I would like to thank Angela, Bob, Christoph, Francesca, Irina, Karen, Lucia, Nandita, Natasha and Sante for many enjoyable hours throughout the course of my study.

## TABLE OF CONTENTS

| CHAPTER |                                                                                      | Page |
|---------|--------------------------------------------------------------------------------------|------|
| I       | INTRODUCTION . . . . .                                                               | 1    |
| II      | TYPE IIA SUPERGRAVITY ON $S^m \times T^n$ . . . . .                                  | 7    |
|         | A. The $S^4$ reduction of eleven-dimensional supergravity . . . .                    | 7    |
|         | B. Type IIA on $S^3$ . . . . .                                                       | 13   |
|         | C. $S^4$ reduction of type IIA supergravity . . . . .                                | 21   |
|         | D. Type IIA supergravity on $S^1 \times S^3$ and $S^1 \times S^3 \times S^1$ . . . . | 25   |
| III     | TYPE IIB SUPERGRAVITY ON $S^m \times T^n$ . . . . .                                  | 27   |
|         | A. Five-dimensional Romans' theory from type IIB on $S^5$ . .                        | 27   |
|         | B. $SO(6)$ gauged supergravity in $D = 5$ . . . . .                                  | 32   |
| IV      | SOME APPLICATIONS . . . . .                                                          | 38   |
|         | A. Super Yang-Mills operators via AdS/CFT correspondence .                           | 38   |
|         | B. Supergravity duals of $\mathcal{N} = 2$ SCFTs in $D = 3$ and $D = 5$ .            | 43   |
|         | 1. Supergravity duals of three-dimensional $\mathcal{N} = 2$ SCFT                    | 44   |
|         | 2. A non-Abelian solution in massive type IIA theory . .                             | 52   |
|         | 3. A non-Abelian solution . . . . .                                                  | 55   |
|         | 4. A black hole solution in the massless theory . . . . .                            | 57   |
|         | 5. A solution with excited B-fields . . . . .                                        | 58   |
|         | C. Supergravity duals of three-dimensional $\mathcal{N} = 1$ SYM                     |      |
|         | theory on a torus . . . . .                                                          | 61   |
|         | 1. Supergravity duals of $\mathcal{N} = 1$ SYM theory in $D = 3$                     |      |
|         | on a torus . . . . .                                                                 | 62   |
|         | 2. Up-lifting to type IIB theory . . . . .                                           | 67   |
|         | 3. Up-lifting to type IIA theory . . . . .                                           | 68   |
|         | 4. The singular $SO(4)$ -symmetric solution . . . . .                                | 69   |
| V       | CONCLUSIONS . . . . .                                                                | 70   |
|         | REFERENCES . . . . .                                                                 | 73   |
|         | APPENDIX A . . . . .                                                                 | 87   |

|                      | Page |
|----------------------|------|
| APPENDIX B . . . . . | 89   |
| APPENDIX C . . . . . | 91   |
| APPENDIX D . . . . . | 93   |
| APPENDIX E . . . . . | 94   |
| APPENDIX F . . . . . | 100  |
| VITA . . . . .       | 104  |

## LIST OF TABLES

| TABLE |                                                                                                                                                | Page |
|-------|------------------------------------------------------------------------------------------------------------------------------------------------|------|
| I     | Structure of ten-dimensional metric . . . . .                                                                                                  | 63   |
| II    | On-shell degrees of freedom in $D$ dimensions. $\alpha = \frac{1}{2}D$ if $D$ is<br>even, $\alpha = \frac{1}{2}(D - 1)$ if $D$ is odd. . . . . | 88   |



## LIST OF FIGURES

| FIGURE |                                                                                                                                                                                                                                                                                                                               | Page |
|--------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 1      | Orbits for the case $AdS_4 \times H_2$ . In the UV limit $F$ behaves as $u^2$ leading to an $AdS_6$ -type region. On the other hand, it flows to “bad” singularities (BS) in the IR. We have used $g = 3m$ with $m = \sqrt{2}$ , and $a = 1$ . . . . .                                                                        | 51   |
| 2      | We show the behavior of the orbits for the case $S^2$ . Similar comments as in the previous case are also valid here. For small $u$ -values we can see the region corresponding to $F \approx \frac{1}{u}$ . . . . .                                                                                                          | 52   |
| 3      | Behavior of the orbits for the non-Abelian massive case. The $AdS_6$ -type region is at the UV, i.e. for large $u$ and $F$ -values. In the IR it flows to “bad” singularities (BS). The cross indicates the fixed point-solution aforementioned. We again set $g = 3m$ , $m = \sqrt{2}$ , and $a = b = \lambda = 1$ . . . . . | 55   |

## CHAPTER I

### INTRODUCTION

Finding an ultimate quantum field theory that describes a unification of four known, fundamental interactions, namely electromagnetic, gravitational, strong and weak interactions, has been a dream for many generations of physicists, and it might still be a dream for many others yet to come. Over the years, there have been many attempts to achieve this dream. Presently, ten-dimensional string theory and eleven-dimensional M-theory [1, 2], whose low-energy limits are ten-dimensional and eleven-dimensional supergravities [3, 4] respectively, are our best candidates for the final theory. To describe our four-dimensional world, the extra dimensions of string theory and M-theory are assumed to be very small and compact, à la the Kaluza-Klein idea [5]. By small, we mean that it is of the order of the Planck scale. One reason the Kaluza-Klein idea is very attractive is that it unifies gravitational and gauge interactions. A satisfactory by-product of learning how to perform dimensional reduction via Kaluza-Klein reduction is that we find many of the lower-dimensional theories we wish to consider are derivable from a single and unique theory, *the Theory of Everything*.

In the 1980s, many different supergravity theories in various dimensions were constructed [6]. Ungauged maximal supergravities in various dimensions from  $D = 10$  to  $D = 3$  can be obtained by Kaluza-Klein reductions of eleven-dimensional supergravity on a  $T^n$  ( $1 \leq n \leq 8$ ) [7]. Reducing these ungauged maximal supergravities on  $T^{1,m}$  torus, which includes a time-like direction, gives rise to Euclidean supergravities in various dimensions [8, 9]. It has long been believed that many gauged supergravities

---

The journal model of this dissertation is Nuclear Physics B.

arise from the Kaluza-Klein reduction of eleven-dimensional supergravity or the type IIA and IIB supergravities on certain spherical internal spaces. It was shown long ago that at the level of linearized fluctuations, the massless spectrum of the maximal gauged supergravities in  $D = 4$  and  $D = 7$  can be obtained from appropriate truncations of those coming from the  $S^7$  and  $S^4$  reductions of  $D = 11$  supergravity [10, 11]. Likewise, the spectrum of maximal gauged supergravity in  $D = 5$  is known to arise from a truncation of the  $S^5$  reduction from type IIB supergravity [12, 13]. Although these linearized results are rather easily established, it is much harder to determine whether the truncations to the massless supermultiplets are consistent at the full non-linear level. The key point of concern here is that once the non-linear interactions are included, the possibility exists that in a full, untruncated reduction, source terms built purely from the fields of the massless multiplet might arise in the field equations for the lower-dimensional massive multiplets, making it inconsistent to set the massive fields to zero. For the Kaluza-Klein reductions on  $S^1$ , and  $n$ -dimensional torus  $T^n$ , which can be viewed as a sequences of  $S^1$  reductions, the truncations to the massless sector are always consistent. The reason for this consistency can be understood in the following way: The massless fields are singlets under the  $U(1)^n$  isometry group of the  $n$ -torus, while the massive modes are non-singlets. The products of singlets under the group action can never generate non-singlets, so there are no couplings between massless and massive modes. Thus, no matter how non-linear the higher-dimensional theory, the Kaluza-Klein reduction on  $T^n$  will be a consistent one. However, the story is significantly different when one performs the Kaluza-Klein reductions on arbitrary spaces, and in particular, on spherical spaces. Indeed, it is not hard to establish that in a spherical reduction of a generic, higher-dimensional theory, there will definitely be such couplings, making a consistent truncation to the massless sector impossible. Thus, if the spherical reductions (with truncation to the massless supermultiplet) in

supergravities are to be consistent, it must be because of remarkable special properties of these particular theories. General arguments suggesting that reductions to the supergravity multiplet should always be consistent were constructed in [14]. Despite all the apparent obstacles, it was shown [15] that the four-dimensional,  $\mathcal{N} = 8$  gauged  $SO(8)$  supergravity can be embedded at the full non-linear level in eleven-dimensional supergravity. However, the result is rather implicit, so practically speaking, it is hard to use. Furthermore, there is no demonstration at the non-linear level for the consistency of the complete  $S^5$  reduction from type IIB supergravity. Therefore, finding explicit, non-linear reduction ansätze remains a challenging problem.

A few years ago, a new duality was conjectured [16, 17, 18], which relates supergravities in anti-de Sitter backgrounds to conformal field theories on their boundaries (AdS/CFT). Since the anti-de Sitter backgrounds arise as solutions of lower-dimensional gauged supergravities, the conjectured AdS/CFT correspondence has led to a revival of interest in deriving these gauged supergravities by Kaluza-Klein reduction from the ten-dimensional fundamental string theories and eleven-dimensional M-theory.

In the last few years, much progress has been achieved in understanding the full non-linear structure of certain Kaluza-Klein sphere reductions. The starting point of the whole program of finding the non-linear reduction ansätze was an observation that the rotating D3-brane should have a “decoupling limit” similar to that of the non-rotating D3-brane.<sup>1</sup> After taking the decoupling limit, the rotating D3-brane metric has the form of an  $AdS_5$  black hole times  $S^5$ . From this observation, we have learned a great deal about the structure of the non-linear reduction ansätze, and we

---

<sup>1</sup>The decoupling limit is a point where the space-time of the non-rotating D3-brane becomes a product space  $M_5 \times S^5$ . If the non-rotating D3-brane is extremal,  $M_5$  is five-dimensional anti-de Sitter space-time ( $AdS_5$ ).

were able to conjecture, and then verify, the correct reduction ansatz. The ansatz describes the exact embedding of the five-dimensional,  $\mathcal{N} = 2$  gauged  $U(1)^3$  supergravity into type IIB supergravity. Similarly, the ansätze which allow the embedding of the  $U(1)^2$  and  $U(1)^4$  truncations of  $D = 7$  and  $D = 4$  gauged supergravities into eleven-dimensional supergravity were constructed [19]. Not long after our work [19], using a formalism based on an analysis of the supersymmetry transformation rules, the complete ansatz for the  $S^4$  reduction of eleven-dimensional supergravity giving rise to the seven-dimensional gauged  $SO(5)$  supergravity was obtained [20, 21]. The consistent reduction of massive type IIA supergravity on a locally  $S^4$  space, to give  $\mathcal{N} = 2$  gauged  $SU(2)$  supergravity in  $D = 6$ , has also been obtained [22]. Further examples of AdS with warped space-times were constructed in [23, 24].

Studying the singular limit of the known  $S^4$  reduction of eleven-dimensional supergravity [21], which gives rise to the standard  $SO(5)$ -gauged maximal supergravity in  $D = 7$ , we constructed a consistent reduction of type IIA on  $S^3$  [25], leading to a maximal gauged supergravity in seven dimensions with the full set of massless  $SO(4)$  Yang-Mills fields. We also obtained the consistent reduction of type IIA supergravity on  $S^4$ , giving rise to an  $SO(5)$ -gauged maximal supergravity in  $D = 6$ . Details of these constructions will be presented in chapter II. Also in chapter II, a consistent reduction ansatz, which allows us to embed any solution of six-dimensional Romans' theory [26] without mass parameter into ten-dimensional supergravity is presented. As will be seen in chapter II, this six-dimensional Romans' theory can be obtained from a reduction of type IIA supergravity on  $S^1 \times S^3$ . In addition, a reduction ansatz of type IIA on  $S^1 \times S^3 \times S^1$ , giving rise to Romans' theory in  $D = 5$  without the  $U(1)$  gauge-coupling, will be also presented in chapter II.

In addition to maximal Abelian truncations in  $D = 4, 5$  and  $7$  [19], certain reductions to truncations of the maximal gauged supergravities in various dimensions

have been constructed. These have the advantage of being considerably simpler than the maximal theories, allowing the reduction ansatz to be presented in a more explicit form. Cases that have been constructed include the  $\mathcal{N} = 2$  gauged  $SU(2)$  supergravity in  $D = 7$  [27]; the  $\mathcal{N} = 4$  gauged  $SU(2) \times U(1)$  in  $D = 5$  [28]; the  $\mathcal{N} = 4$  gauged  $SO(4)$  supergravity in  $D = 4$  [29]. Using these results, certain ungauged supergravities in  $D = 4$ ,  $D = 5$  and  $D = 6$  can be embedded into type IIA, IIB and eleven-dimensional supergravities [30, 31]. One can also consider non-supersymmetric truncations of the maximal gauged supergravities. The consistent truncations of the  $D = 7$ ,  $D = 5$  and  $D = 4$  supergravities to the graviton plus scalar subsectors comprising only the diagonal scalars in the  $SL(N, \mathbb{R})/SO(N)$  scalar sub-manifolds were considered (with  $N = 5, 6$  and  $8$ , respectively), and the consistent embeddings in  $D = 11$  and  $D = 10$  were constructed [32, 33, 34]. Another complete ansatz for the consistent embedding of a subsector, which consists of 20 scalars and 15 Yang-Mills fields, of the five-dimensional  $\mathcal{N} = 8$  gauged  $SO(6)$  supergravity into type IIB was obtained [35]. Various ansätze of type IIB supergravity on  $S^5$  and on  $S^3 \times T^2$  will also be presented in chapter III.

In chapter IV, certain applications of non-linear Kaluza-Klein ansätze will be presented. In particular, we will show [36] that with the full non-linear reduction ansatz the calculation of  $n$ -point correlation functions in super Yang-Mills (SYM) theory via AdS/CFT correspondence appears to be much simpler than those previously appearing in the literature, which use the linearized ansatz followed by non-linear field redefinitions [37].

Recently, non-linear Kaluza-Klein reductions ansätze have been extensively used in finding supergravity solutions corresponding to certain twisted field theories [38]. This searching provides some new examples of AdS/CFT correspondence. Wrapping M5-branes and D3-branes on Riemann-surfaces that are holomorphically embedded

in Calabi-Yau 2- or 3-folds, giving rise to four-dimensional field theories with  $\mathcal{N} = 2, 1$  supersymmetry and two-dimensional field theories with  $(4, 4)$  and  $(2, 2)$  supersymmetry respectively, were analyzed [39]. Type IIB NS5-branes wrapped on  $S^2$  leading to four-dimensional  $\mathcal{N} = 1$  super Yang-Mills theory was studied [40]. Some related work associated with Dp-branes and M5-branes wrapping Riemann surfaces in Calabi-Yau spaces had studied earlier in [41, 23, 42, 43, 44, 45, 46]. These studies have been extended for the cases of M5-branes and NS5-branes wrapping on associative 3-cycles in seven-dimensional manifolds with  $G_2$  holonomy [47]; D3-branes [48] wrapped on associative 3-cycles; M5-branes wrapping on Kähler 4-cycles, special Lagrangian 3-, 4-, and 5-cycles, co-associative 4-cycles and Caley 4-cycles [49]; D4-branes as well as NS5-branes wrapping on 2- and 3-cycles [50]; D6-branes wrapping on  $S^2$  and  $S^3$  [51]; NS5-branes wrapping on  $S^3 \times T^2$  [52, 53]; M2-branes wrapped on 2-cycles in Calabi-Yau 2-, 3-, 4- and 5-folds [54]; D6-branes wrapped on co-associative 4-cycles with constant curvature of seven manifold of  $G_2$  holonomy [55]; 5-branes wrapping on  $S^2$  in a Calabi-Yau 2-folds [56]; D6-branes wrapping on Kähler 4-cycles [57, 58]. We will also present some supergravity solutions [50, 52] which are dual to certain supersymmetric gauge theories via AdS/CFT correspondence in chapter IV.

Chapter V is devoted to conclusions. Brief reviews of eleven-dimensional supergravity and type IIA, IIB and massive type IIA in  $D = 10$  are presented in appendices A, B, C, and D, respectively. Appendix E is devoted to a review of six-dimensional,  $SU(2)$ -gauged supergravity and the  $\mathcal{N} = 4$ ,  $SU(2) \times U(1)$ -gauged supergravity in  $D = 5$  is presented in appendix F.

## CHAPTER II

TYPE IIA SUPERGRAVITY ON  $S^m \times T^n$ A. The  $S^4$  reduction of eleven-dimensional supergravity

The complete ansatz for the  $S^4$  reduction of eleven-dimensional supergravity was obtained in [21], using a formalism based on an analysis of the supersymmetry transformation rules. One may also study the reduction from a purely bosonic standpoint, by verifying that if the ansatz is substituted into the eleven-dimensional equations of motion, it consistently yields the equations of motion of the seven-dimensional gauged  $SO(5)$  supergravity. We shall carry out this procedure here, in order to establish notation, and to obtain the complete system of seven-dimensional bosonic equations of motion, which we shall need in the later part.

After some manipulation, the Kaluza-Klein  $S^4$  reduction ansatz obtained in [21] for eleven-dimensional supergravity can be expressed as follows:

$$d\hat{s}_{11}^2 = \Delta^{1/3} ds_7^2 + \frac{1}{g^2} \Delta^{-2/3} T_{ij}^{-1} \mathcal{D}\mu^i \mathcal{D}\mu^j, \quad (2.1)$$

$$\begin{aligned} \hat{F}_{(4)} = & \frac{1}{4!} \epsilon_{i_1 \dots i_5} \left[ -\frac{1}{g^3} U \Delta^{-2} \mu^{i_1} \mathcal{D}\mu^{i_2} \wedge \dots \wedge \mathcal{D}\mu^{i_5} \right. \\ & + \frac{4}{g^3} \Delta^{-2} T^{i_1 m} \mathcal{D}T^{i_2 n} \mu^m \mu^n \mathcal{D}\mu^{i_3} \wedge \dots \wedge \mathcal{D}\mu^{i_5} \\ & \left. + \frac{6}{g^2} \Delta^{-1} F_{(2)}^{i_1 i_2} \wedge \mathcal{D}\mu^{i_3} \wedge \mathcal{D}\mu^{i_4} T^{i_5 j} \mu^j \right] - T_{ij} * S_{(3)}^i \mu^j + \frac{1}{g} S_{(3)}^i \wedge \mathcal{D}\mu^i, \end{aligned} \quad (2.2)$$

$$\begin{aligned} *\hat{F}_{(4)} = & -gU\epsilon_{(7)} - \frac{1}{g} T_{ij}^{-1} * \mathcal{D}T^{ik} \mu_k \wedge \mathcal{D}\mu^j + \frac{1}{2g^2} T_{ik}^{-1} T_{j\ell}^{-1} * F_{(2)}^{ij} \wedge \mathcal{D}\mu^k \wedge \mathcal{D}\mu^\ell \\ & - \frac{1}{6g^3} \Delta^{-1} \epsilon_{ij\ell_1 \ell_2 \ell_3} * S_{(3)}^m T_{im} T_{jk} \mu^k \wedge \mathcal{D}\mu^{\ell_1} \wedge \mathcal{D}\mu^{\ell_2} \wedge \mathcal{D}\mu^{\ell_3} \\ & + \frac{1}{g^4} \Delta^{-1} T_{ij} S_{(3)}^i \mu^j \wedge W, \end{aligned} \quad (2.3)$$



where

$$\begin{aligned}
U &\equiv 2T_{ij}T_{jk}\mu^i\mu^k - \Delta T_{ii}, \quad \Delta \equiv T_{ij}\mu^i\mu^j, \quad F_{(2)}^{ij} = \mathcal{D}A_{(1)}^{ij} \equiv dA_{(1)}^{ij} + gA_{(1)}^{ik} \wedge A_{(1)}^{kj}, \\
\mathcal{D}\mu^i &\equiv d\mu^i + gA_{(1)}^{ij}\mu^j, \quad \mathcal{D}T_{ij} \equiv dT_{ij} + gA_{(1)}^{ik}T_{kj} + gA_{(1)}^{jk}T_{ik}, \quad \mu^i\mu^i \equiv 1, \\
W &\equiv \frac{1}{24}\epsilon_{i_1\dots i_5}\mu^{i_1}\mathcal{D}\mu^{i_2} \wedge \dots \wedge \mathcal{D}\mu^{i_5},
\end{aligned} \tag{2.4}$$

where the symmetric matrix  $T_{ij}$  is unimodular and hat denotes eleven-dimensional fields. Furthermore,  $T_{ij}$  parameterizes the scalar coset  $SL(6, \mathbb{R})/SO(6)$ .

To obtain seven-dimensional equations of motion, we first consider the Bianchi identity  $d\hat{F}_{(4)} = 0$ . Substituting (2.2) into this, we obtain the following equations:

$$\mathcal{D}(T_{ij}*S_{(3)}^j) = F_{(2)}^{ij} \wedge S_{(3)}^j, \tag{2.5}$$

$$H_{(4)}^i = gT_{ij}*S_{(3)}^j + \frac{1}{8}\epsilon_{ij_1\dots j_4}F_{(2)}^{j_1j_2} \wedge F_{(2)}^{j_3j_4}, \tag{2.6}$$

where we define

$$H_{(4)}^i \equiv \mathcal{D}S_{(3)}^i = dS_{(3)}^i + gA_{(1)}^{ij} \wedge S_{(3)}^j. \tag{2.7}$$

Next, we substitute the ansatz into the  $D = 11$  field equation (A.5) for  $\hat{F}_{(4)}$  and we get

$$\begin{aligned}
\mathcal{D}(T_{ik}^{-1}T_{j\ell}^{-1}*F_{(2)}^{ij}) &= -2gT_{i[k}*\mathcal{D}T_{\ell]i} - \frac{1}{2g}\epsilon_{i_1i_2i_3k\ell}F_2^{i_1i_2}H_{(4)}^{i_3} \\
&\quad + \frac{3}{2g}\delta_{i_1i_2k\ell}^{j_1j_2j_3j_4}F_{(2)}^{i_1i_2} \wedge F_{(2)}^{j_1j_2} \wedge F_{(2)}^{j_3j_4} - S_{(3)}^k \wedge S_{(3)}^\ell.
\end{aligned} \tag{2.8}$$

$$\begin{aligned}
\mathcal{D}(T_{ik}^{-1}*\mathcal{D}T_{kj}) &= 2g^2(2T_{ik}T_{kj} - T_{kk}T_{ij})\epsilon_{(7)} + T_{im}^{-1}T_{k\ell}^{-1}*F_{(2)}^{m\ell} \wedge F_{(2)}^{kj} \\
&\quad + T_{jk}*S_{(3)}^k \wedge S_{(3)}^i - \frac{1}{5}\delta_{ij}\left[2g^2(2T_{ik}T_{ik} - 2(T_{ii})^2)\epsilon_{(7)}\right. \\
&\quad \left.+ T_{nm}^{-1}T_{k\ell}^{-1}*F_{(2)}^{m\ell} \wedge F_{(2)}^{kn} + T_{k\ell}*S_{(3)}^k \wedge S_{(3)}^\ell\right],
\end{aligned} \tag{2.9}$$

for the Yang-Mills and scalar equations of motion in  $D = 7$ . Note from (2.8) that it would be inconsistent to set the Yang-Mills fields to zero while retaining the scalars

$T_{ij}$ , since the currents  $T_{i[k}^{-1} * \mathcal{D}T_{\ell]i}$  act as sources for them. A truncation where the Yang-Mills fields are set to zero is consistent, however, if the scalars are also truncated to the diagonal subsector  $T_{ij} = \text{diag}(X_1, X_2, \dots, X_6)$ , as in the consistent reductions constructed in [32, 33].

We find that all the equations of motion can be derived from the following seven-dimensional Lagrangian

$$\begin{aligned} \mathcal{L}_7 = & R * \mathbb{1} - \frac{1}{4} T_{ij}^{-1} * \mathcal{D}T_{jk} \wedge T_{k\ell}^{-1} \mathcal{D}T_{\ell i} - \frac{1}{4} T_{ik}^{-1} T_{j\ell}^{-1} * F_{(2)}^{ij} \wedge F_{(2)}^{k\ell} - \frac{1}{2} T_{ij} * S_{(3)}^i \wedge S_{(3)}^j \\ & + \frac{1}{2g} S_{(3)}^i \wedge H_{(4)}^i - \frac{1}{8g} \epsilon_{ij_1 \dots j_4} S_{(3)}^i \wedge F_{(2)}^{j_1 j_2} \wedge F_{(2)}^{j_3 j_4} + \frac{1}{g} \Omega_{(7)} - V * \mathbb{1} , \end{aligned} \quad (2.10)$$

where  $H_{(4)}^i$  are given by (2.7) and the potential  $V$  is given by

$$V = \frac{1}{2} g^2 (2T_{ij} T_{ij} - (T_{ii})^2) , \quad (2.11)$$

and  $\Omega_{(7)}$  is a Chern-Simons type of term built from the Yang-Mills fields, which has the property that its variation with respect to  $A_{(1)}^{ij}$  gives

$$\delta \Omega_{(7)} = \frac{3}{4} \delta_{i_1 i_2 k \ell}^{j_1 j_2 j_3 j_4} F_{(2)}^{i_1 i_2} \wedge F_{(2)}^{j_1 j_2} \wedge F_{(2)}^{j_3 j_4} \wedge \delta A_{(1)}^{k \ell} . \quad (2.12)$$

Note that the  $S_{(3)}^i$  are viewed as fundamental fields in the Lagrangian, and that (2.6) is their first-order equation. In fact (2.10) is precisely the bosonic sector of the Lagrangian describing maximal gauged seven-dimensional supergravity that was derived in [59]. An explicit expression for the 7-form  $\Omega_{(7)}$  can be found there.

Although we have fully checked the eleven-dimensional Bianchi identity and field equation for  $\hat{F}_{(4)}$  here, we have not completed the task of substituting the ansatz into the eleven-dimensional Einstein equations. This would be an extremely complicated calculation, on account of the Yang-Mills gauge fields. However, various complete consistency checks, including the higher-dimensional Einstein equation, have been

performed in various truncations of the full  $\mathcal{N} = 4$  maximal supergravity embedding, including the  $\mathcal{N} = 2$  gauged theory in [27], and the non-supersymmetric truncation in [33] where the gauge fields are set to zero and only the diagonal scalars in  $T_{ij}$  are retained. All the evidence points to the full consistency of the reduction.<sup>1</sup>

It is perhaps worth making a few further remarks on the nature of the reduction ansatz. One might wonder whether the ansatz (2.2) on the 4-form field strength  $\hat{F}_{(4)}$  could be re-expressed as an ansatz on its potential  $\hat{A}_{(3)}$ . As it stands, (2.2) only satisfies the Bianchi identity  $d\hat{F}_{(4)} = 0$  by virtue of the lower-dimensional equations (2.5) and (2.6). However, if (2.6) is substituted into (2.2), we obtain an expression that satisfies  $d\hat{F}_{(4)} = 0$  without the use of any lower-dimensional equations. However, one does still have to make use of the fact that the  $\mu^i$  coordinates satisfy the constraint  $\mu^i \mu^i = 1$ , and this prevents one from writing an explicit ansatz for  $\hat{A}_{(3)}$  that has a manifest  $SO(5)$  symmetry. One could solve for one of the  $\mu^i$  in terms of the others, but this would break the manifest local symmetry from  $SO(5)$  to  $SO(4)$ . In principle though, this could be done, and then one could presumably substitute the resulting ansatz directly into the eleven-dimensional Lagrangian. After integrating out the internal 4-sphere directions, one could then in principle obtain a seven-dimensional Lagrangian in which, after re-organizing terms, the local  $SO(5)$  symmetry could again become manifest.

It should, of course, be emphasized that merely substituting an ansatz into a Lagrangian and integrating out the internal directions to obtain a lower-dimensional Lagrangian is justifiable only if one already has an independent proof of the consis-

---

<sup>1</sup>The original demonstration in [21], based on the reduction of the eleven-dimensional supersymmetry transformation rules, also provides extremely compelling evidence. Strictly speaking, the arguments presented there also fall short of a complete and rigorous proof, since they involve an approximation in which the quartic fermion terms in the theory are neglected.

tency of the proposed reduction ansatz.<sup>2</sup> If one is in any case going to work with the higher-dimensional field equations in order to prove the consistency, it is not clear that there would be any significant benefit to be derived from then re-expressing the ansatz in a form where it could be substituted into the Lagrangian.

It is interesting to observe that one cannot take the limit  $g \rightarrow 0$  in the Lagrangian (2.10), on account of the terms proportional to  $g^{-1}$  in the second line. We know, on the other hand, that it must be possible to recover the ungauged  $D = 7$  theory by turning off the gauge coupling constant. In fact the problem is associated with a pathology in taking the limit at the level of the Lagrangian, rather than in the equations of motion. This can be seen by looking instead at the seven-dimensional equations of motion, which were given earlier. The only apparent obstacle to taking the limit  $g \rightarrow 0$  is in the Yang-Mills equations (2.8), but in fact this is illusory. If we substitute the first-order equation (2.6) into (2.8) it gives

$$\mathcal{D}(T_{ik}^{-1}T_{j\ell}^{-1}*F_{(2)}^{ij}) = -2gT_{i[k}^{-1}*\mathcal{D}T_{\ell]i} - \frac{1}{2}\epsilon_{i_1i_2i_3k\ell}F_2^{i_1i_2} \wedge T_{ij}*S_{(3)}^j - S_{(3)}^k \wedge S_{(3)}^\ell, \quad (2.13)$$

which has a perfectly smooth  $g \rightarrow 0$  limit. It is clear that equations of motion (2.6) and (2.9) and the Einstein equations of motion also have a smooth limit. (The reason why the Einstein equations have the smooth limit is because the  $1/g$  terms in the Lagrangian (2.10) do not involve the metric, and thus they give no contribution.)

Unlike in the gauged theory, we should not treat the  $S_{(3)}^i$  fields as fundamental variables in a Lagrangian formulation in the ungauged limit. This is because once the gauge coupling  $g$  is sent to zero, the fields  $S_{(3)}^i$  behave like 3-form field strengths. This

---

<sup>2</sup>A classic illustration is provided by the example of five-dimensional pure gravity with an (inconsistent) Kaluza-Klein reduction in which the scalar dilaton is omitted. Substituting this into the five-dimensional Einstein-Hilbert action yields the perfectly self-consistent Einstein-Maxwell action in  $D = 4$ , but fails to reveal that setting the scalar to zero is inconsistent with the internal component of the five-dimensional Einstein equation.

can be seen from the first-order equation of motion (2.6), which in the limit  $g \rightarrow 0$  becomes

$$dS_{(3)}^i = \frac{1}{8} \epsilon_{ij_1 \dots j_4} dA_{(1)}^{j_1 j_2} \wedge dA_{(1)}^{j_3 j_4} , \quad (2.14)$$

and should now be interpreted as a Bianchi identity. This can be solved by introducing 2-form gauge potentials  $A_{(2)}^i$ , with the  $S_{(3)}^i$  given by

$$S_{(3)}^i = dA_{(2)}^i + \frac{1}{8} \epsilon_{ij_1 \dots j_4} A_{(1)}^{j_1 j_2} \wedge dA_{(1)}^{j_3 j_4} . \quad (2.15)$$

In terms of these 2-form potentials, the equations of motion can now be obtained from the Lagrangian

$$\begin{aligned} \mathcal{L}_7^0 = & R * \mathbf{1} - \frac{1}{4} T_{ij}^{-1} * dT_{jk} \wedge T_{k\ell}^{-1} dT_{\ell i} - \frac{1}{4} T_{ik}^{-1} T_{j\ell}^{-1} * F_{(2)}^{ij} \wedge F_{(2)}^{k\ell} - \frac{1}{2} T_{ij} * S_{(3)}^i \wedge S_{(3)}^j \\ & + \frac{1}{2} A_{(1)}^{ij} \wedge S_{(3)}^i \wedge S_{(3)}^j - 2S_{(3)}^i \wedge A_{(2)}^j \wedge dA_{(1)}^{ij} , \end{aligned} \quad (2.16)$$

where  $S_{(3)}^i$  is given by (2.15). This is precisely the bosonic Lagrangian of the ungauged maximal supergravity in  $D = 7$ .

It is worth exploring in a little more detail why it is possible to take a smooth  $g \rightarrow 0$  limit in the seven-dimensional equations of motion, but not in the Lagrangian. We note that in this limit the Lagrangian (2.10) can be expressed as

$$\mathcal{L}_7 = \frac{1}{g} L + \mathcal{O}(1) , \quad (2.17)$$

where

$$L = \frac{1}{2} S_{(3)}^i dS_{(3)}^i - \frac{1}{8} \epsilon_{ij_1 \dots j_4} S_{(3)}^i \wedge F_{(2)}^{j_1 j_2} \wedge F_{(2)}^{j_3 j_4} + \Omega_{(7)} . \quad (2.18)$$

The term  $L/g$ , which diverges in the  $g \rightarrow 0$  limit, clearly emphasizes that the Lagrangian (2.16) is not merely the  $g \rightarrow 0$  limit of (2.10). However if we make use of the equations of motion, we find that in the  $g \rightarrow 0$  limit the  $S_{(3)}^i$  can be solved by

(2.15). Substituting this into (2.18), we find that in this limit it becomes

$$L = \frac{1}{16} \epsilon_{ij_1 \dots j_4} dA_{(2)}^i \wedge dA_{(1)}^{j_1 j_2} \wedge dA_{(1)}^{j_3 j_4} + \mathcal{O}(g) , \quad (2.19)$$

and so the singular terms in  $L/g$  form a total derivative and hence can be subtracted from the Lagrangian. This analysis explains why it is possible to take a smooth  $g \rightarrow 0$  limit in the equations of motion, but not in the Lagrangian.

### B. Type IIA on $S^3$

Here we examine, at the level of the seven-dimensional theory itself, how to take a limit in which the  $SO(5)$ -gauged sector is broken down to  $SO(4)$ . In a later section, we shall show how this can be interpreted as an  $S^3$  reduction of type IIA supergravity. We shall do that by showing how to take a limit in which the internal  $S^4$  in the original reduction from  $D = 11$  becomes  $\mathbb{R} \times S^3$ . For now, however, we shall examine the  $SO(4)$ -gauged limit entirely from the perspective of the seven-dimensional theory itself.

To take the limit, we break the  $SO(5)$  covariance by splitting the  $\underline{5}$  index  $i$  as

$$i = (0, \alpha) , \quad (2.20)$$

where  $1 \leq \alpha \leq 4$ . We also introduce a constant parameter  $\lambda$ , which will be sent to zero as the limit is taken. We find that the various seven-dimensional fields, and the  $SO(5)$  gauge-coupling constant, should be scaled as follows:

$$\begin{aligned} g &= \lambda^2 \tilde{g} , & A_{(1)}^{0\alpha} &= \lambda^3 \tilde{A}_{(1)}^{0\alpha} , & A_{(1)}^{\alpha\beta} &= \lambda^{-2} \tilde{A}_{(1)}^{\alpha\beta} , \\ S_{(3)}^0 &= \lambda^{-4} \tilde{S}_{(3)}^0 , & S_{(3)}^\alpha &= \lambda \tilde{S}_{(3)}^\alpha . \end{aligned} \quad (2.21)$$

$$T_{ij}^{-1} = \begin{pmatrix} \lambda^{-8} \Phi & \lambda^{-3} \Phi \chi_\alpha \\ \lambda^{-3} \Phi \chi_\alpha & \lambda^2 M_{\alpha\beta}^{-1} + \lambda^2 \Phi \chi_\alpha \chi_\beta \end{pmatrix} .$$

As we show in the next section, this rescaling corresponds to a degeneration of  $S^4$  to  $R \times S^3$ . Note that in this rescaling, we have also performed a decomposition of the scalar matrix  $T_{ij}^{-1}$  that is of the form of a Kaluza-Klein metric decomposition. It is useful also to present the consequent decomposition for  $T_{ij}$ , which turns out to be

$$T_{ij} = \begin{pmatrix} \lambda^8 \Phi^{-1} + \lambda^8 \chi_\gamma \chi^\gamma & -\lambda^3 \chi^\alpha \\ -\lambda^3 \chi^\alpha & \lambda^{-2} M_{\alpha\beta} \end{pmatrix} . \quad (2.22)$$

Calculating the determinant, we get

$$\det(T_{ij}) = \Phi^{-1} \det(M_{\alpha\beta}) . \quad (2.23)$$

Since we know that  $\det(T_{ij}) = 1$ , it follows that

$$\Phi = \det(M_{\alpha\beta}) . \quad (2.24)$$

The fields  $\chi_\alpha$  are “axionic” scalars. Note that we shall also have

$$\begin{aligned} H_{(4)}^0 &= \lambda^{-4} \widetilde{H}_{(4)}^0 , & H_{(4)}^\alpha &= \lambda \widetilde{H}_{(4)}^\alpha , \\ \widetilde{H}_{(4)}^0 &= d\widetilde{S}_{(3)}^0 , & \widetilde{H}_{(4)}^\alpha &= \widetilde{\mathcal{D}}\widetilde{S}_{(3)}^\alpha - \tilde{g} \widetilde{A}_{(1)}^{0\alpha} \wedge \widetilde{S}_{(3)}^0 . \end{aligned} \quad (2.25)$$

We have defined an  $SO(4)$ -covariant exterior derivative  $\widetilde{\mathcal{D}}$ , which acts on quantities with  $SO(4)$  indexes  $\alpha, \beta, \dots$  in the obvious way:

$$\widetilde{\mathcal{D}} X_\alpha = dX_\alpha + \tilde{g} \widetilde{A}_{(1)}^{a\beta} X_\beta , \quad (2.26)$$

etc. It is helpful also to make the following further field redefinitions:

$$\begin{aligned} G_{(2)}^\alpha &\equiv \widetilde{F}_{(2)}^{0\alpha} + \chi_\beta \widetilde{F}_{(2)}^{\beta\alpha} , \\ G_{(3)}^\alpha &\equiv \widetilde{S}_{(3)}^\alpha - \chi_\alpha \widetilde{S}_{(3)}^0 , \end{aligned} \quad (2.27)$$

$$G_{(1)}^\alpha \equiv \tilde{\mathcal{D}}\chi_\alpha - \tilde{g}\tilde{A}_{(1)}^{0\alpha},$$

where  $\tilde{F}_{(2)}^{0\alpha} \equiv \tilde{\mathcal{D}}\tilde{A}_{(1)}^{0\alpha}$ .

We may now substitute these redefined fields into the seven-dimensional equations of motion. We find that a smooth limit in which  $\lambda$  is sent to zero exists, leading to an  $SO(4)$ -gauged seven-dimensional theory. Our results for the seven-dimensional equations of motion are as follows. The fields  $H_{(4)}^i$  become

$$\tilde{H}_{(4)}^0 = d\tilde{S}_{(3)}^0, \quad \tilde{H}_{(4)}^\alpha = \tilde{\mathcal{D}}G_{(3)}^\alpha + G_{(1)}^\alpha \wedge \tilde{S}_{(3)}^0 + \chi_\alpha dS_{(3)}^0. \quad (2.28)$$

The first-order equations (2.7) give

$$\begin{aligned} \tilde{H}_{(4)}^0 &= \frac{1}{8}\epsilon_{\alpha_1\cdots\alpha_4}\tilde{F}_{(2)}^{\alpha_1\alpha_2}\wedge\tilde{F}_{(2)}^{\alpha_3\alpha_4}, \\ \tilde{F}_{(4)}^\alpha &= \tilde{g}M_{\alpha\beta}*G_{(3)}^\beta - \frac{1}{2}\epsilon_{\alpha\beta\gamma\delta}G_{(2)}^\beta\wedge\tilde{F}_{(2)}^{\gamma\delta} - G_{(1)}^\alpha\wedge\tilde{S}_{(3)}^0, \end{aligned} \quad (2.29)$$

where we have defined

$$F_{(4)}^\alpha \equiv \tilde{\mathcal{D}}G_{(3)}^\alpha. \quad (2.30)$$

The second-order equations (2.5) (which are nothing but Bianchi identities following from (2.7)) become

$$\begin{aligned} d(\Phi^{-1}*\tilde{S}_{(3)}^0) &= M_{\alpha\beta}*G_{(3)}^\alpha\wedge G_{(1)}^\beta + G_{(2)}^\alpha\wedge G_{(3)}^\alpha, \\ \tilde{\mathcal{D}}(M_{\alpha\beta}*G_{(3)}^\beta) &= \tilde{F}_{(2)}^{\alpha\beta}\wedge G_{(3)}^\beta - G_{(2)}^\alpha\wedge\tilde{S}_{(3)}^0. \end{aligned} \quad (2.31)$$

The Yang-Mills equations (2.8) become

$$\begin{aligned} \tilde{\mathcal{D}}(\Phi M_{\alpha\beta}^{-1}*G_{(2)}^\beta) &= \tilde{g}\Phi M_{\alpha\beta}*G_{(1)}^\beta - \tilde{S}_{(3)}^0\wedge G_{(3)}^\alpha - \frac{1}{2}\epsilon_{\alpha\beta_1\beta_2\beta_3}M_{\beta_3\gamma}\tilde{F}_{(2)}^{\beta_1\beta_2}\wedge*G_{(3)}^\gamma, \\ \tilde{\mathcal{D}}[M_{\gamma\alpha}^{-1}M_{\delta\beta}^{-1}*\tilde{F}_{(2)}^{\gamma\delta}] &= -2\tilde{g}M_{\gamma[\alpha}^{-1}*\tilde{\mathcal{D}}M_{\beta]\gamma} - G_{(3)}^\alpha\wedge G_{(3)}^\beta + \Phi M_{\alpha\gamma}^{-1}G_{(1)}^\beta\wedge*G_{(2)}^\gamma \\ &\quad - \Phi M_{\beta\gamma}^{-1}G_{(1)}^\alpha\wedge*G_{(2)}^\gamma - \epsilon_{\alpha\beta\gamma\delta}M_{\delta\lambda}G_{(2)}^\gamma\wedge*G_{(3)}^\lambda \\ &\quad - \frac{1}{2}\Phi^{-1}\epsilon_{\alpha\beta\gamma\delta}\tilde{F}_{(2)}^{\gamma\delta}\wedge*\tilde{S}_{(3)}^0. \end{aligned} \quad (2.32)$$



Finally, the scalar field equations (2.9) give the following:

$$\begin{aligned}
d(\Phi^{-1} * d\Phi) &= \Phi M_{\alpha\beta} * G_1^\alpha \wedge G_{(1)}^\beta + \Phi M_{\alpha\beta}^{-1} * G_{(2)}^\alpha \wedge G_{(2)}^\beta \\
&\quad - \Phi^{-1} * \tilde{S}_{(3)}^0 \wedge \tilde{S}_{(3)}^0 + \frac{1}{5} Q , \\
\tilde{\mathcal{D}}(\Phi M_{\alpha\beta} * G_{(1)}^\beta) &= \Phi M_{\beta\gamma}^{-1} * G_{(2)}^\gamma \wedge \tilde{F}_{(2)}^{\alpha\beta} - M_{\alpha\beta} * G_{(3)}^\beta \wedge \tilde{S}_{(3)}^0 , \\
\tilde{\mathcal{D}}(M_{\alpha\gamma}^{-1} * \tilde{\mathcal{D}} M_{\gamma\beta}) &= \Phi M_{\beta\gamma} * G_{(1)}^\gamma \wedge G_{(1)}^\alpha + M_{\beta\gamma} * G_{(3)}^\gamma \wedge G_{(3)}^\alpha \\
&\quad - \Phi M_{\alpha\gamma}^{-1} * G_{(2)}^\gamma \wedge G_{(2)}^\beta + M_{\alpha\gamma}^{-1} M_{\lambda\delta}^{-1} * \tilde{F}_{(2)}^{\gamma\delta} \wedge \tilde{F}_{(2)}^{\lambda\beta} \\
&\quad + 2\tilde{g}^2 (2M_{\alpha\gamma} M_{\gamma\beta} - M_{\gamma\gamma} M_{\alpha\beta}) \epsilon_{(\tau)} - \frac{1}{5} \delta_{\alpha\beta} Q . \quad (2.33)
\end{aligned}$$

In these equations, the quantity  $Q$  is the limit of the trace term multiplying  $\delta_{ij}$  in (2.9), and is given by

$$\begin{aligned}
Q &= 2\tilde{g}^2 (2M_{\alpha\beta} M_{\alpha\beta} - (M_{\alpha\alpha})^2) \epsilon_{(\tau)} - M_{\alpha\gamma}^{-1} M_{\beta\delta}^{-1} * \tilde{F}_{(2)}^{\alpha\beta} \wedge \tilde{F}_{(2)}^{\gamma\delta} \\
&\quad + \Phi^{-1} * \tilde{S}_{(3)}^0 \wedge \tilde{S}_{(3)}^0 - 2\Phi M_{\alpha\beta}^{-1} * G_{(2)}^\alpha \wedge G_{(2)}^\beta + M_{\alpha\beta} * G_{(3)}^\alpha \wedge G_{(3)}^\beta . \quad (2.34)
\end{aligned}$$

Having obtained the seven-dimensional equations of motion for the  $SO(4)$ -gauged limit, we can now seek a Lagrangian from which they can be generated. A crucial point is that the equations involving  $\tilde{H}_{(4)}^0$  in (2.28) and (2.29) give

$$d\tilde{S}_{(3)}^0 = \frac{1}{8} \epsilon_{\alpha_1 \dots \alpha_4} \tilde{F}_{(2)}^{\alpha_1 \alpha_2} \wedge \tilde{F}_{(2)}^{\alpha_3 \alpha_4} , \quad (2.35)$$

which allows us to strip off the exterior derivative by writing

$$\tilde{S}_{(3)}^0 = dA_{(2)} + \omega_{(3)} , \quad (2.36)$$

where  $\tilde{S}_{(3)}^0$  is now viewed as a field strength with 2-form potential  $A_{(2)}$ , and

$$\omega_{(3)} \equiv \frac{1}{8} \epsilon_{\alpha_1 \dots \alpha_4} (\tilde{F}_{(2)}^{\alpha_1 \alpha_2} \wedge \tilde{A}_{(1)}^{\alpha_3 \alpha_4} - \frac{1}{3} \tilde{g} \tilde{A}_{(1)}^{\alpha_1 \alpha_2} \wedge \tilde{A}_{(1)}^{\alpha_3 \beta} \wedge \tilde{A}_{(1)}^{\beta \alpha_4}) . \quad (2.37)$$

We can now see that the equations of motion can be derived from the following seven-

dimensional Lagrangian, in which  $A_{(2)}$ , and not its field strength  $\tilde{S}_{(3)}^0 \equiv dA_{(2)} + \omega_{(3)}$ , is viewed as a fundamental field:

$$\begin{aligned}
\mathcal{L}_7 = & R * \mathbb{1} - \frac{1}{4} \Phi^{-2} * d\Phi \wedge d\Phi - \frac{1}{4} \tilde{M}_{\alpha\beta}^{-1} * \tilde{\mathcal{D}} \tilde{M}_{\beta\gamma} \wedge \tilde{M}_{\gamma\delta}^{-1} \tilde{\mathcal{D}} \tilde{M}_{\delta\alpha} - \frac{1}{2} \Phi^{-1} * \tilde{S}_{(3)}^0 \wedge \tilde{S}_{(3)}^0 \\
& - \frac{1}{4} M_{\alpha\gamma}^{-1} M_{\beta\delta}^{-1} * \tilde{F}_{(2)}^{\alpha\beta} \wedge \tilde{F}_{(2)}^{\gamma\delta} - \frac{1}{2} \Phi M_{\alpha\beta}^{-1} * G_{(2)}^\alpha \wedge G_{(2)}^\beta - \frac{1}{2} \Phi M_{\alpha\beta} * G_{(1)}^\alpha \wedge G_{(1)}^\beta \\
& - \frac{1}{2} M_{\alpha\beta} * G_{(3)}^\alpha \wedge G_{(3)}^\beta - \tilde{V} * \mathbb{1} + \frac{1}{2\tilde{g}} \tilde{\mathcal{D}} \tilde{S}_{(3)}^\alpha \wedge \tilde{S}_{(3)}^\alpha + \tilde{S}_{(3)}^\alpha \wedge \tilde{S}_{(3)}^0 \wedge A_{(1)}^{0\alpha} + \frac{1}{\tilde{g}} \tilde{\Omega}_{(7)} \\
& + \frac{1}{2\tilde{g}} \epsilon_{\alpha\beta\gamma\delta} \tilde{S}_{(3)}^\alpha \wedge \tilde{F}_{(2)}^{0\beta} \wedge \tilde{F}_{(2)}^{\gamma\delta} + \frac{1}{4} \epsilon_{\alpha_1 \dots \alpha_4} \tilde{S}_{(3)}^0 \wedge \tilde{F}_{(2)}^{\alpha_1 \alpha_2} \wedge \tilde{A}_{(1)}^{0\alpha_3} \wedge \tilde{A}_{(1)}^{0\alpha_4}, \quad (2.38)
\end{aligned}$$

where  $\tilde{\Omega}_{(7)}$  is built purely from  $\tilde{A}_{(1)}^{\alpha\beta}$  and  $\tilde{A}_{(1)}^{0\alpha}$ . It is defined by the requirement that its variations with respect to  $\tilde{A}_{(1)}^{\alpha\beta}$  and  $\tilde{A}_{(1)}^{0\alpha}$  should produce the necessary terms in the equations of motion for these fields. Since it has a rather complicated structure, we shall not present it here. Note that  $\tilde{M}_{\alpha\beta} \equiv \Phi^{-1/4} M_{\alpha\beta}$ , where  $\tilde{M}_{\alpha\beta}$  is the unimodular matrix.

Using the above scaling limit of the gauged  $SO(5)$  theory in seven dimensions, in which an  $SO(4)$  gauging survives, we show that it leads to a degeneration in which the 4-sphere becomes  $\mathbb{R} \times S^3$ . We can then re-interpret the reduction from  $D = 11$  as an initial “ordinary” Kaluza-Klein reduction step from  $D = 11$  to give the type IIA supergravity in  $D = 10$ , followed by a non-trivial reduction of the type IIA theory on  $S^3$ , in which the entire  $SO(4)$  isometry group is gauged. Note that the  $S^3$  reduction of type IIA supergravity discussed in [60], giving a seven-dimensional theory with just an  $SU(2)$  gauging, was re-derived in [61] as a singular limit of the  $S^4$  reduction of  $D = 11$  supergravity that was obtained in [21]. Since the  $S^3$  reduction in [60] retains only the left-acting  $SU(2)$  of the  $SO(4) \sim SU(2)_L \times SU(2)_R$  of gauge fields, the consistency of that reduction is guaranteed by group-theoretic arguments, based on the fact that all the retained fields are singlets under the right-acting  $SU(2)_R$ . The subtleties of the consistency of the  $S^4$  reduction in [21] are therefore lost in the

singular limit to  $\mathbb{R} \times S^3$  discussed in [61], since a truncation to the  $SU(2)_L$  subgroup of the  $SO(4)$  gauge group is made. By contrast, the  $\mathbb{R} \times S^3$  singular limit that we consider here retains all the fields of the  $S^4$  reduction in [21], and the proof of the consistency of the resulting  $S^3$  reduction of the type IIA theory follows from the non-trivial consistency of the reduction in [21], and has no simple group-theoretic explanation.

To take this limit, we combine the scalings of seven-dimensional quantities derived in the previous section with an appropriately-matched rescaling of the coordinates  $\mu^i$  defined on the internal 4-sphere. As in [62], we see that after splitting the  $\mu^i$  into  $\mu^0$  and  $\mu^\alpha$ , these additional scalings should take the form

$$\mu^0 = \lambda^5 \tilde{\mu}^0, \quad \mu^\alpha = \tilde{\mu}^\alpha. \quad (2.39)$$

In the limit where  $\lambda$  goes to zero, we see that the original constraint  $\mu^i \mu^i = 1$  becomes

$$\tilde{\mu}^\alpha \tilde{\mu}^\alpha = 1, \quad (2.40)$$

implying that the  $\tilde{\mu}^\alpha$  coordinates define a 3-sphere, while the coordinate  $\tilde{\mu}^0$  is now unconstrained and ranges over the real line  $\mathbb{R}$ .

Combining this with the rescalings of the previous section, we find that the  $S^4$  metric reduction ansatz (2.1) becomes

$$\begin{aligned} d\hat{s}_{11}^2 = & \lambda^{-2/3} \left[ \tilde{\Delta}^{1/3} ds_7^2 + \frac{1}{\tilde{g}^2} \tilde{\Delta}^{-2/3} M_{\alpha\beta}^{-1} \tilde{\mathcal{D}}\tilde{\mu}^\alpha \tilde{\mathcal{D}}\tilde{\mu}^\beta \right. \\ & \left. + \frac{1}{\tilde{g}^2} \tilde{\Delta}^{-2/3} \Phi (d\tilde{\mu}_0 + \tilde{g} \tilde{A}_{(1)}^{0\alpha} \tilde{\mu}^\alpha + \chi_\alpha \tilde{\mathcal{D}}\tilde{\mu}^\alpha)^2 \right], \end{aligned} \quad (2.41)$$

where

$$\tilde{\Delta} \equiv M_{\alpha\beta} \tilde{\mu}^\alpha \tilde{\mu}^\beta. \quad (2.42)$$

Thus  $\tilde{\mu}_0$  can be interpreted as the “extra” coordinate of a standard type of Kaluza-

Klein reduction from  $D = 11$  to  $D = 10$ , with

$$d\hat{s}_{11}^2 = e^{-\frac{1}{6}\phi} ds_{10}^2 + e^{\frac{4}{3}\phi} (d\tilde{\mu}_0 + \mathcal{A}_{(1)})^2 . \quad (2.43)$$

By comparing (2.43) with (2.41), we can read off the  $S^3$  reduction ansatz for the ten-dimensional fields. Thus we find that the ten-dimensional metric is reduced according to

$$ds_{10}^2 = \Phi^{1/8} \left[ \tilde{\Delta}^{1/4} ds_7^2 + \frac{1}{\tilde{g}^2} \tilde{\Delta}^{-3/4} M_{\alpha\beta}^{-1} \tilde{\mathcal{D}}\tilde{\mu}^\alpha \tilde{\mathcal{D}}\tilde{\mu}^\beta \right] , \quad (2.44)$$

while the ansatz for the dilaton  $\phi$  of the ten-dimensional theory is

$$e^{2\phi} = \tilde{\Delta}^{-1} \Phi^{3/2} . \quad (2.45)$$

Finally, the reduction ansatz for the ten-dimensional Kaluza-Klein vector is

$$\mathcal{A}_{(1)} = \tilde{g} \tilde{A}_{(1)}^{0\alpha} \tilde{\mu}^\alpha + \chi_\alpha \tilde{\mathcal{D}}\tilde{\mu}^\alpha . \quad (2.46)$$

These results for the  $S^3$  reduction of the ten-dimensional metric and dilaton agree precisely with the results obtained in [63]. (Note that the field  $\Phi$  is called  $Y$  there, and our  $M_{\alpha\beta}$  is called  $T_{ij}$  there.) Note that the field strength  $\mathcal{F}_{(2)} = d\mathcal{A}_{(1)}$  following from (2.46) has the simple expression

$$\mathcal{F}_{(2)} = \tilde{g} G_{(2)}^\alpha \tilde{\mu}^\alpha + G_{(1)}^\alpha \wedge \tilde{\mathcal{D}}\tilde{\mu}^\alpha . \quad (2.47)$$

So far, we have read off the reduction ansätze for those fields of ten-dimensional type IIA supergravity that come from the reduction of the eleven-dimensional metric. The remaining type IIA fields come from the reduction of the eleven-dimensional 4-form. Under the standard Kaluza-Klein procedure, this reduces as follows:

$$\hat{F}_{(4)} = F_{(4)} + F_{(3)} \wedge (d\tilde{\mu}^0 + \mathcal{A}_{(1)}) . \quad (2.48)$$

By applying the  $\lambda$ -rescaling derived previously to the  $S^4$  reduction ansatz (2.2) for the eleven-dimensional 4-form, and comparing with (2.48), we obtain the following expressions for the  $S^3$  reduction ansätze for the ten-dimensional 4-form and 3-form fields:

$$F_{(4)} = \frac{\tilde{\Delta}^{-1}}{\tilde{g}^3} M_{\alpha\beta} G_{(1)}^\alpha \tilde{\mu}^\beta \wedge \tilde{W} + \frac{\tilde{\Delta}^{-1}}{2\tilde{g}^2} \epsilon_{\alpha_1 \dots \alpha_4} M_{\alpha_4\beta} \tilde{\mu}^\beta G_{(2)}^{\alpha_1} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_2} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_3} \\ - M_{\alpha\beta} * G_{(3)}^\alpha \tilde{\mu}^\beta + \frac{1}{\tilde{g}} G_{(3)}^\alpha \wedge \tilde{\mathcal{D}}\tilde{\mu}^\alpha, \quad (2.49)$$

$$F_{(3)} = -\frac{\tilde{U}\tilde{\Delta}^{-2}}{\tilde{g}^3} \tilde{W} + \frac{\tilde{\Delta}^{-2}}{2\tilde{g}^3} \epsilon_{\alpha_1 \dots \alpha_4} M_{\alpha_1\beta} \tilde{\mu}^\beta \tilde{\mathcal{D}}M_{\alpha_2\gamma} \tilde{\mu}^\gamma \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_3} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_4} \\ + \frac{\tilde{\Delta}^{-1}}{2\tilde{g}^2} \epsilon_{\alpha_1 \dots \alpha_4} M_{\alpha_1\beta} \tilde{\mu}^\beta \tilde{F}_{(2)}^{\alpha_2\alpha_3} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_4} + \frac{1}{\tilde{g}} \tilde{S}_{(3)}^0, \quad (2.50)$$

where

$$\tilde{W} \equiv \frac{1}{6} \epsilon_{\alpha_1 \dots \alpha_4} \tilde{\mu}^{\alpha_1} \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_2} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_3} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_4}. \quad (2.51)$$

It is also useful to present the expressions for the ten-dimensional Hodge duals of the field strengths:

$$e^{\frac{3}{2}\phi} * \mathcal{F}_{(2)} = \frac{\tilde{\Delta}^{-1}\Phi}{\tilde{g}^5} * G_{(2)}^\alpha \tilde{\mu}^\alpha \wedge \tilde{W} + \frac{\tilde{\Delta}^{-1}\Phi}{2\tilde{g}^4} \epsilon_{\alpha_1 \dots \alpha_4} M_{\alpha_1\beta} \tilde{\mu}^\beta M_{\alpha_2\gamma} * G_{(1)}^\gamma \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_3} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_4}, \\ e^{-\phi} * F_{(3)} = -\tilde{g}\tilde{U}\epsilon_{(7)} - \frac{1}{\tilde{g}^3} \Phi^{-1} * \tilde{S}_{(3)}^0 \wedge \tilde{W} \\ + \frac{1}{2\tilde{g}^2} M_{\alpha\gamma}^{-1} M_{\beta\delta}^{-1} * \tilde{F}_{(2)}^{\alpha\beta} \wedge \tilde{\mathcal{D}}\tilde{\mu}^\gamma \wedge \tilde{\mathcal{D}}\tilde{\mu}^\delta - \frac{1}{\tilde{g}} M_{\alpha\beta}^{-1} * \tilde{\mathcal{D}}M_{\alpha\gamma} \tilde{\mu}^\gamma \wedge \tilde{\mathcal{D}}\tilde{\mu}^\beta, \\ e^{\frac{1}{2}\phi} * F_{(4)} = \frac{1}{\tilde{g}} \Phi M_{\alpha\beta} * G_{(1)}^\alpha \tilde{\mu}^\beta - \frac{1}{\tilde{g}^2} \Phi M_{\alpha\beta}^{-1} * G_{(2)}^\alpha \wedge \tilde{\mathcal{D}}\tilde{\mu}^\beta + \frac{\tilde{\Delta}^{-1}}{\tilde{g}^4} M_{\alpha\beta} G_{(3)}^\alpha \tilde{\mu}^\beta \wedge \tilde{W} \\ + \frac{\tilde{\Delta}^{-1}}{2\tilde{g}^3} \epsilon_{\alpha_1 \dots \alpha_4} M_{\alpha_1\beta} \tilde{\mu}^\beta M_{\alpha_2\gamma} * G_{(3)}^\gamma \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_3} \wedge \tilde{\mathcal{D}}\tilde{\mu}^{\alpha_4}. \quad (2.52)$$

(Here we are using  $*$  to denote a Hodge dualization in the ten-dimensional metric  $ds_{10}^2$ , to distinguish it from  $*$  which denotes the seven-dimensional Hodge dual in the metric  $ds_7^2$ .)

The consistency of the  $S^3$  reduction of the type IIA theory using the ansatz that

we obtained in the previous subsection is guaranteed by virtue of the consistency of the  $S^4$  reduction from  $D = 11$ . It is still useful, however, to examine the reduction directly, by substituting the ansatz into the equations of motion of type IIA supergravity (B.5). By this means we can obtain an explicit verification of the validity of the limiting procedures that we applied in obtaining the  $S^3$  reduction ansatz. Note that it is consistent to truncate the theory to the NS-NS sector, namely the subsector comprising the metric, the dilaton and the 3-form field strength. This implies that it is possible also to perform an  $S^3$  reduction of the NS-NS sector alone, which was indeed demonstrated in [63]. On the other hand it is not consistent to truncate the theory to a sub-sector comprising only the metric, the dilaton and the 4-form field strength, which again is in agreement with the conclusion in [63] that it is not consistent to perform an  $S^4$  reduction on such a subsector. However, as we show in the next section, there is a consistent  $S^4$  reduction if we include all the fields of the type IIA theory.

### C. $S^4$ reduction of type IIA supergravity

In this section, we derive the ansatz for the consistent  $S^4$  reduction of type IIA supergravity from the  $S^4$  reduction ansatz of eleven-dimensional supergravity. In this case we do not need to take any singular limit of the internal 4-sphere, but rather, we extract the “extra” coordinate from the seven-dimensional space-time of the original eleven-dimensional supergravity reduction ansatz. The resulting six-dimensional  $SO(5)$ -gauged maximal supergravity can be obtained from the Kaluza-Klein reduction of seven-dimensional gauged maximal supergravity on a circle.

We begin, therefore, by making a standard  $S^1$  Kaluza-Klein reduction of the

seven-dimensional metric:

$$ds_7^2 = e^{-2\alpha\varphi} ds_6^2 + e^{8\alpha\varphi} (dz + \bar{\mathcal{A}}_{(1)})^2, \quad (2.53)$$

where  $\alpha = 1/\sqrt{40}$ . With this parameterization the metric reduction preserves the Einstein frame, and the dilatonic scalar  $\varphi$  has the canonical normalization for its kinetic term in six dimensions.<sup>3</sup> Substituting (2.53) into the original metric reduction ansatz (2.1), we obtain

$$d\hat{s}_{11}^2 = \Delta^{1/3} e^{-2\alpha\varphi} ds_6^2 + \frac{1}{g^2} \Delta^{-2/3} T_{ij}^{-1} \mathcal{D}\mu^i \mathcal{D}\mu^j + \Delta^{1/3} e^{8\alpha\varphi} (dz + \bar{\mathcal{A}}_{(1)})^2. \quad (2.54)$$

In order to extract the ansatz for the  $S^4$  reduction of type IIA supergravity, we must first rewrite (2.54) in the form

$$d\hat{s}_{11}^2 = e^{-\frac{1}{6}\phi} ds_{10}^2 + e^{\frac{4}{3}\phi} (dz + \mathcal{A}_{(1)})^2, \quad (2.55)$$

which is a canonical  $S^1$  reduction from  $D = 11$  to  $D = 10$ . It is not immediately obvious that this can easily be done, since the Yang-Mills fields  $A_{(1)}^{ij}$  appearing in the covariant differentials  $\mathcal{D}\mu^i$  in (2.54) must themselves be reduced according to standard Kaluza-Klein rules,

$$A_{(1)}^{ij} = \bar{A}_{(1)}^{ij} + \chi^{ij} (dz + \bar{\mathcal{A}}_{(1)}), \quad (2.56)$$

where  $\bar{A}_{(1)}^{ij}$  are the  $SO(5)$  gauge potentials in six dimensions, and  $\chi^{ij}$  are axions in  $D = 6$ . Thus we have

$$\mathcal{D}\mu^i = \bar{\mathcal{D}}\mu^i + g \chi^{ij} \mu^j (dz + \bar{\mathcal{A}}_{(1)}), \quad (2.57)$$

---

<sup>3</sup>We use a bar to denote six-dimensional fields, in cases where this is necessary to avoid an ambiguity.

where

$$\bar{\mathcal{D}}\mu^i \equiv d\mu^i + g \bar{A}_{(1)}^{ij} \mu^j . \quad (2.58)$$

This means that the differential  $dz$  actually appears in a much more complicated way in (2.54) than is apparent at first sight. Nonetheless, we find that one can in fact “miraculously” complete the square, and thereby rewrite (2.54) in the form of (2.55).

To present the result, it is useful to make the following definitions:

$$\begin{aligned} \Omega &\equiv \Delta^{1/3} e^{8\alpha\varphi} + \Delta^{-2/3} T_{ij}^{-1} \chi^{ik} \chi^{j\ell} \mu^k \mu^\ell , \\ Z_{ij} &\equiv T_{ij}^{-1} - \Omega^{-1} \Delta^{-2/3} T_{ik}^{-1} T_{j\ell}^{-1} \chi^{km} \chi^{\ell n} \mu^m \mu^n . \end{aligned} \quad (2.59)$$

In terms of these, we find after some algebra that we can rewrite (2.54) as

$$d\hat{s}_{11}^2 = \Delta^{1/3} e^{-2\alpha\varphi} ds_6^2 + \frac{1}{g^2} \Delta^{-2/3} Z_{ij} \bar{\mathcal{D}}\mu^i \bar{\mathcal{D}}\mu^j + \Omega (dz + \mathcal{A}_{(1)})^2 , \quad (2.60)$$

where the ten-dimensional potential  $\mathcal{A}_{(1)}$  is given in terms of six-dimensional fields by

$$\mathcal{A}_{(1)} = \bar{\mathcal{A}}_{(1)} + \frac{1}{g} \Omega^{-1} \Delta^{-2/3} T_{ij}^{-1} \chi^{jk} \mu^k \bar{\mathcal{D}}\mu^i . \quad (2.61)$$

This is therefore the Kaluza-Klein  $S^4$  reduction ansatz for the 1-form  $\mathcal{A}_{(1)}$  of the type IIA theory. Comparing (2.60) with (2.55), we see that the Kaluza-Klein reduction ansätze for the metric  $ds_{10}^2$  and dilaton  $\phi$  of the type IIA theory are given by

$$\begin{aligned} ds_{10}^2 &= \Omega^{1/8} \Delta^{1/3} e^{-2\alpha\varphi} ds_6^2 + \frac{1}{g^2} \Omega^{1/8} \Delta^{-2/3} Z_{ij} \bar{\mathcal{D}}\mu^i \bar{\mathcal{D}}\mu^j , \\ e^{\frac{4}{3}\phi} &= \Omega . \end{aligned} \quad (2.62)$$

The  $S^4$  reduction ansatz for the R-R 4-form  $F_{(4)}$  of the type IIA theory is obtained in a similar manner, by first implementing a standard  $S^1$  Kaluza-Klein reduction on the various seven-dimensional fields appearing in the  $S^4$  reduction ansatz (2.2) for the eleven-dimensional 4-form  $\hat{F}_{(4)}$ , and then matching this to a standard  $S^1$  reduction of



$\hat{F}_{(4)}$  from  $D = 11$  to  $D = 10$ :

$$\hat{F}_{(4)} = F_{(4)} + F_{(3)} \wedge (dz + \mathcal{A}_{(1)}) . \quad (2.63)$$

Note that in doing this, it is appropriate to treat the 3-form fields  $S_{(3)}^i$  of the seven-dimensional theory as field strengths for the purpose of the  $S^1$  reduction to  $D = 6$ , viz.

$$S_{(3)}^i = \bar{S}_{(3)}^i + \bar{S}_{(2)}^i \wedge (dz + \bar{\mathcal{A}}_{(1)}) . \quad (2.64)$$

It is worth noting also that this implies that the reduction of the seven-dimensional Hodge duals  $*S_{(3)}^i$  will be given by

$$*S_{(3)}^i = e^{4\alpha\varphi} \bar{*}\bar{S}_{(3)}^i \wedge (dz + \bar{\mathcal{A}}_{(1)}) + e^{-6\alpha\varphi} \bar{*}\bar{S}_{(2)}^i , \quad (2.65)$$

where  $\bar{*}$  denotes a Hodge dualization in the six-dimensional metric  $ds_6^2$ .

With these preliminaries, it is now a mechanical, albeit somewhat uninspiring, exercise to make the necessary substitutions into (2.2), and, by comparing with (2.63), read off the expressions for  $F_{(4)}$  and  $F_{(3)}$ . These give the Kaluza-Klein  $S^4$  reductions ansätze for the 4-form and 3-form field strengths of type IIA supergravity. We shall not present the results explicitly here, since they are rather complicated, and are easily written down “by inspection” if required. For these purposes, the following identities are useful:

$$\begin{aligned} (dz + \bar{\mathcal{A}}_{(1)}) &= (dz + \mathcal{A}_{(1)}) - \frac{1}{g} \Omega^{-1} \Delta^{-2/3} T_{ij}^{-1} \chi^{jk} \mu^k \tilde{\mathcal{D}}\mu^i , \\ \mathcal{D}X_i &= \bar{\mathcal{D}}X_i - \Omega^{-1} \Delta^{-2/3} \chi^{ij} X_j T_{k\ell}^{-1} \chi^{\ell m} \mu^m \bar{\mathcal{D}}\mu^k + g \chi^{ij} (dz + \mathcal{A}_{(1)}) , \\ \mathcal{D}\mu^i &= T_{ij} Z_{jk} \bar{\mathcal{D}}\mu^k + g \chi^{ij} \mu^j (dz + \mathcal{A}_{(1)}) , \end{aligned} \quad (2.66)$$

where in the last line  $X_i$  represents any six-dimensional field in the vector representation of  $SO(5)$ , and the covariant derivative generalizes to higher-rank  $SO(5)$  tensors

in the obvious way.

If we substitute the  $S^4$  reduction ansätze given for the ten-dimensional dilaton, metric and 1-form in (2.62), and (2.61), together with those for  $F_{(4)}$  and  $F_{(3)}$  as described above, into the equations of motion of type IIA supergravity, we shall obtain a consistent reduction to six dimensions. This six-dimensional theory will be precisely the one that follows by performing an ordinary  $S^1$  Kaluza-Klein reduction on the  $SO(5)$ -gauged maximal supergravity in  $D = 7$ , whose bosonic Lagrangian is given in (2.38).

It is perhaps worth remarking that the expression (2.62) for the Kaluza-Klein  $S^4$  reduction of the type IIA supergravity metric illustrates a point that has been observed previously (for example in [29, 34]), namely that the ansatz becomes much more complicated when axions or pseudoscalars are involved. Although the axions  $\chi^{ij}$  would not be seen in the metric ansatz in a linearized analysis, they make an appearance in a rather complicated way in the full non-linear ansatz that we have obtained here, for example in the quantities  $\Omega$  and  $Z_{ij}$  defined in (2.59). They will also, of course, appear in the ansätze for the  $F_{(4)}$  and  $F_{(3)}$  field strengths. It may be that the results we are finding here could be useful in other contexts, for providing clues as to how the axionic scalars should appear in the Kaluza-Klein reduction ansatz.

#### D. Type IIA supergravity on $S^1 \times S^3$ and $S^1 \times S^3 \times S^1$

As shown in [22] and reviewed in appendix E, the six-dimensional  $SU(2)$ -gauged supergravity can be derived from reducing massive type IIA on a local  $S^4$ . In this section, we present two other reduction ansätze which lead to two subsectors of Romans' theories in  $D = 5$  and  $D = 6$ .

Firstly, the dual six-dimensional Romans' theory (E.14) can be obtained from a

subset, which consists of a graviton, a dilaton and a 4-form field strength, of type IIA supergravity on  $S^1 \times S^3$ . The idea is that we first reduce type IIA on  $S^1$ . After consistently setting 4-form and one scalar to zero, we then reduce the obtained nine-dimensional theory on  $S^3$  using the formulae in [63]. The reduction ansätze are

$$\begin{aligned} d\hat{s}_{10}^2 &= -\frac{1}{2}e^{\frac{\sqrt{2}}{4}\phi} ds_6^2 + \frac{1}{g^2}e^{-\frac{3\sqrt{2}}{4}\phi} \sum_{i=1}^3 \left( \sigma^i - \frac{1}{\sqrt{2}}g A_{(1)}^i \right)^2 + e^{\frac{5\sqrt{2}}{4}\phi} dZ^2, \\ \hat{F}_{(4)} &= \left( F_{(3)} - \frac{1}{g^2}h^1 \wedge h^2 \wedge h^3 + \frac{1}{\sqrt{2}}g F_{(2)}^i \wedge h^i \right) \wedge dZ, \\ \hat{\phi} &= \frac{1}{\sqrt{2}}\phi, \end{aligned} \tag{2.67}$$

where  $h^i = \sigma^i - \frac{1}{\sqrt{2}}g A^i$ . As pointed out in [50], the equations of motion produced by the ansätze (2.67) are precisely those of the  $\mathcal{N} = \tilde{4}^g$  Romans' theory (E.14).

After dualizing 3-form field following (E.16), we obtain the Romans' theory in  $D = 6$  (Appendix E). Reducing this theory on  $S^1$  produces the Romans' theory in  $D = 5$  without  $U(1)$  gauge coupling (Appendix F). This implies that the five-dimensional Romans' theory without the  $U(1)$  gauge-coupling can also be embedded into type IIA supergravity. The reduction ansätze for metric, dilaton and field strength are

$$\begin{aligned} d\hat{s}_{10}^2 &= e^{\frac{7}{8\sqrt{6}}\phi} ds_5^2 + \frac{1}{4g^2}e^{-\frac{9}{8\sqrt{6}}\phi} \sum_{i=1}^3 \left( \sigma^i - g A_1^i \right)^2 + e^{\frac{15}{8\sqrt{6}}\phi} dY^2 + e^{-\frac{9}{8\sqrt{6}}\phi} dZ^2, \\ \hat{F}_4 &= \left( e^{\frac{4}{\sqrt{6}}\phi} * \mathcal{F}_2 - \frac{1}{24g^2} \epsilon_{ijk} h^i \wedge h^j \wedge h^k + \frac{1}{4g} F_2^i \wedge h^i \right) \wedge dY, \\ \hat{\phi} &= \frac{3}{4\sqrt{6}}\phi. \end{aligned} \tag{2.68}$$

## CHAPTER III

TYPE IIB SUPERGRAVITY ON  $S^m \times T^n$ A. Five-dimensional Romans' theory from type IIB on  $S^5$ 

To construct the reduction ansätze, we follow a procedure similar to that used in [27, 22]. We take as our starting point the  $U(1)^3$  Abelian truncation obtained in [19], which involved two independent scalar fields. One can perform a consistent truncation of this, in which two of the three  $U(1)$  potentials are set equal, and at the same time one of the scalar degrees of freedom is eliminated. After doing so, the metric on the internal 5-sphere takes the following form

$$d\omega_5^2 = X \Delta d\xi^2 + X^2 s^2 (d\tau - g B_{(1)})^2 + \frac{1}{4} X^{-1} c^2 \left( d\theta^2 + \sin^2 \theta d\phi^2 + (d\psi + \cos \theta d\phi - \sqrt{2} g A_{(1)})^2 \right), \quad (3.1)$$

where  $X \equiv e^{-\frac{1}{\sqrt{6}}\phi}$ , and we have defined

$$c \equiv \cos \xi, \quad s \equiv \sin \xi. \quad (3.2)$$

The function  $\Delta$  is given by

$$\Delta = X^{-2} s^2 + X c^2. \quad (3.3)$$

At  $X = 1$ , in the absence of the  $U(1)$  gauge fields, the metric (3.1) describes a unit 5-sphere. (The gauge field  $A_{(1)}$  appearing in (3.1) is the one that comes from setting two of the original  $U(1)$  gauge fields in [19] equal.)

We can now make a non-Abelian generalization of the the 5-sphere metric ansatz, by introducing the three left-invariant 1-forms  $\sigma^i$  on the 3-sphere, which satisfy the

Cartan-Maurer equation

$$d\sigma^i + \frac{1}{2}\epsilon_{ijk}\sigma^j \wedge \sigma^k = 0. \quad (3.4)$$

These can be written in terms of the Euler angles as

$$\begin{aligned} \sigma_1 &= \cos\psi d\theta + \sin\theta \sin\psi d\phi, \\ \sigma_2 &= -\sin\psi d\theta + \sin\theta \cos\psi d\phi, \\ \sigma_3 &= d\psi + \cos\theta d\phi. \end{aligned} \quad (3.5)$$

Thus we are naturally led to generalize (3.1) to

$$d\omega_5^2 = X \Delta d\xi^2 + X^2 s^2 (d\tau - gB_{(1)})^2 + \frac{1}{4}X^{-1}c^2 \sum_{i=1}^3 (\sigma^i - \sqrt{2}g A_{(1)}^i)^2. \quad (3.6)$$

The Abelian limit (3.1) is recovered by setting the  $i = 1$  and  $i = 2$  components of the  $SU(2)$  gauge potentials to zero.

By proceeding along these lines, we are eventually led to the following ansätze for the ten-dimensional metric, the 5-form field strength  $\hat{H}_{(5)} \equiv \hat{G}_{(5)} + \hat{*}\hat{G}_{(5)}$ , and the two 2-form potentials:

$$\begin{aligned} ds_{10}^2 &= \Delta^{1/2} ds_5^2 + g^{-2} X \Delta^{1/2} d\xi^2 + g^{-2} \Delta^{-1/2} X^2 s^2 (d\tau - g B_{(1)})^2 \\ &\quad + \frac{1}{4} g^{-2} \Delta^{-1/2} X^{-1} c^2 \sum_i (\sigma^i - \sqrt{2}g A_{(1)}^i)^2, \\ \hat{G}_{(5)} &= 2g U \epsilon_5 - \frac{3sc}{g} X^{-1} *dX \wedge d\xi + \frac{c^2}{8\sqrt{2}g^2} X^{-2} *F_{(2)}^i \wedge h^j \wedge h^k \epsilon_{ijk} \\ &\quad - \frac{sc}{2\sqrt{2}g^2} X^{-2} *F_{(2)}^i \wedge h^i \wedge d\xi - \frac{sc}{g^2} X^4 *G_{(2)} \wedge d\xi \wedge (d\tau - gB_{(1)}), \\ \hat{A}_{(2)} &\equiv \hat{A}_{(2)}^1 + i \hat{A}_{(2)}^2 = -\frac{s}{\sqrt{2}g} e^{-i\tau} A_{(2)}, \\ \hat{\phi} &= 0, \quad \hat{\chi} = 0, \end{aligned} \quad (3.7)$$

where  $h^i \equiv \sigma^i - \sqrt{2}g A_{(1)}^i$ ,  $U \equiv X^2 c^2 + X^{-1} s^2 + X^{-1}$ , and  $\epsilon_5$  is the volume form in the five-dimensional space-time metric  $ds_5^2$ . Note that we have defined the complex 2-

form potential  $\hat{A}_{(2)} \equiv \hat{A}_{(2)}^1 + i \hat{A}_{(2)}^2$  in the type IIB theory. The ten-dimensional dilaton and the axion are constants, which without loss of generality we have set to zero. The  $SU(2)$  Yang-Mills field strengths  $F_{(2)}^i$  are given by  $F_{(2)}^i = dA_{(1)}^i + \frac{1}{\sqrt{2}} g \epsilon_{ijk} A_{(1)}^j \wedge A_{(1)}^k$ . It should be emphasized that all the fields  $X$ ,  $A_{(2)}^\alpha$  and  $A_{(1)}^i$ , and the metric  $ds_5^2$ , appearing on the right-hand sides of (3.7), are taken to depend only on the coordinates of the five-dimensional space-time.<sup>1</sup> Note also that since the two 2-form potentials of the five-dimensional supergravity are charged under the  $U(1)$  factor in the  $SU(2) \times U(1)$  gauge group, these fields appear in the reduction ansatz with the appropriate  $\tau$  dependence for charged  $U(1)$  harmonics.

The ten-dimensional Hodge dual of  $\hat{G}_{(5)}$  turns out to be

$$\begin{aligned} \star \hat{G}_{(5)} = & -\frac{sc^3}{4g^4} U \Delta^{-2} d\xi \wedge \omega \wedge \epsilon_3 + \frac{3s^2c^4}{8g^4} X^{-2} \Delta^{-2} dX \wedge \omega \wedge \epsilon_3 \\ & -\frac{s^2c^2}{8\sqrt{2}g^3} X^{-2} \Delta^{-1} F_{(2)}^i \wedge h^j \wedge h^k \wedge \omega \epsilon_{ijk} + \frac{sc}{2\sqrt{2}g^3} F_{(2)}^i \wedge h^i \wedge d\xi \wedge \omega \\ & -\frac{c^4}{8g^3} X \Delta^{-1} G_{(2)} \wedge \epsilon_3, \end{aligned} \quad (3.8)$$

where  $\omega \equiv d\tau - gB_{(1)}$ , and  $\epsilon_3 \equiv h^1 \wedge h^2 \wedge h^3$ .

We must now verify that the ansätze (3.7) do indeed yield a consistent reduction of the type IIB theory to the  $\mathcal{N} = 4$  gauged supergravity in  $D = 5$ . Let us begin by noting that the Bianchi identity for  $\hat{H}_{(5)}$ , and the field equations for the NS-NS and

---

<sup>1</sup>Note that the metric reduction ansatz in (3.7) has a fairly simple form, and fits in with the general pattern of Kaluza-Klein metric ansätze, such as have been seen in many previous examples. (See [64, 65, 66] for some earlier examples, and [67, 19, 27, 22] for some more recent ones.) It is the determination of the reduction ansätze for the field strengths that is the more difficult part of the problem, and since the consistency of the reduction depends crucially on conspiracies between the field strength and metric contributions, it is only when the full ansatz is known that the consistency can be established.

R-R 3-forms of the type IIB theory become, in the complex notation,

$$d\hat{H}_{(5)} = \frac{i}{2} \bar{\hat{F}}_{(3)} \wedge \hat{F}_{(3)} , \quad d\hat{*}\hat{F}_{(3)} = -i \hat{H}_{(5)} \wedge \hat{F}_{(3)} . \quad (3.9)$$

We now find that the Bianchi identity for  $\hat{H}_{(5)}$  gives rise to the following equations in  $D = 5$ :

$$\begin{aligned} d(X^{-1} * dX) &= \frac{4}{3} g^2 (X^{-1} - X^2) * \mathbb{1} - \frac{1}{6} X^{-2} (*F_{(2)}^i \wedge F_{(2)}^i + *\bar{A}_{(2)} \wedge A_{(2)}) \\ &\quad + \frac{1}{3} X^4 *G_{(2)} \wedge G_{(2)} , \\ d(X^{-2} *F_{(2)}^i) &= \sqrt{2} g X^{-2} \epsilon^{ijk} *F_{(2)}^j \wedge A_{(1)}^k - F_{(2)}^i \wedge G_{(2)} , \\ d(X^4 *G_{(2)}) &= -\frac{1}{2} F_{(2)}^i \wedge F_{(2)}^i - \frac{1}{2} \bar{A}_{(2)} \wedge A_{(2)} . \end{aligned} \quad (3.10)$$

These are precisely equations of motion of the Romans' theory.

Plugging the ansätze (3.7) into the equation of motion for  $\hat{F}_{(3)}$  in (3.9) gives rise to the five-dimensional first-order equation for  $A_{(2)}$  in (F.14), and also the second-order equation

$$\mathcal{D}(X^2 *F_{(3)}) = g^2 X^{-2} *A_{(2)} , \quad (3.11)$$

where the  $U(1)$  gauge-covariant exterior derivative  $\mathcal{D}$  is as defined in (F.15). This second order equation of motion is in fact implied by the first order equation. With our ansätze, the ten-dimensional equations of motion for the dilaton and axion are automatically satisfied. Substituting the ansätze into the ten-dimensional Einstein equation (C.3) does not lead to any other equations except five-dimensional Einstein equation (F.14). Thus the full consistency of the reduction ansatz (3.7) is established.

From the above ansätze, one can perform some appropriate rescaling to transform them from one gauge  $g$  coupling to two gauge couplings  $g_1$  and  $g_2$  which correspond to the gauge couplings of  $U(1)$  and  $SU(2)$  respectively. We then in principle can take a singular limit in which  $g_1$  is set to zero. In this singular limit  $S^5$  is deformed to  $S^3 \times T^2$ .

As a result of this process, we obtain the five-dimensional Romans' theory without the  $U(1)$  gauge-coupling. Instead of taking the singular limit of the metric and field strength ansätze in Eq. (3.7), we obtain the same theory by reducing a subset, which comprises a metric, a dilaton and a NS-NS 3-form, of type IIB supergravity on  $S^3 \times T^2$  together with appropriate dualizations [52]. The ansätze are

$$\begin{aligned}
d\hat{s}_{10}^2 &= e^{\frac{13}{5\sqrt{6}}\phi} ds_5^2 + e^{\frac{3}{5\sqrt{6}}\phi} (dY^2 + dZ^2) + \frac{1}{4g^2} e^{-\frac{3}{\sqrt{6}}\phi} \sum_{i=1}^3 (\sigma^i - g A_1^i)^2, \\
\hat{F}_3 &= e^{\frac{4}{\sqrt{6}}\phi} * \mathcal{F}_2 - \frac{1}{24g^2} \epsilon_{ijk} h^i \wedge h^j \wedge h^k + \frac{1}{4} \sum_{i=1}^3 F_2^i \wedge h^i, \\
\hat{\phi} &= \sqrt{6} \phi, \quad \text{and} \quad h^i = \sigma^i - g A_1^i.
\end{aligned} \tag{3.12}$$

Notice that reducing the above mentioned subset of type IIB supergravity produces the  $SU(2)$ -gauged  $\mathcal{N} = 2$  supergravity in  $D = 7$  without topological mass term [68, 69]. Reductions of seven-dimensional gauged supergravity on  $S^1$  and on  $T^2$  were done in [70, 71, 72]. The above mentioned seven-dimensional gauged supergravity can also be obtained from appropriate truncation of a gauged supergravity derived from reducing type IIA supergravity on  $S^3$  [25]. It is not clear that a supergravity obtained from reducing type IIA on  $S^3$  is dual to a theory obtained from reducing IIB on  $S^3$  in the same sense as T-duality (sometimes it is called non-Abelian T-duality). However, reducing particular subsets of type IIA and type IIB supergravities on  $S^3$  produces the same theory. Based on this one might say that there exists non-Abelian T-duality. Unfortunately, an explicit construction from purely supergravity theories has yet been done. Arguments based on worldsheet conformal field theory were done by many groups [73, 74, 75, 76, 77, 78, 79].



### B. $SO(6)$ gauged supergravity in $D = 5$

In this section, we obtain the consistent reduction ansatz for a different truncation of the maximal gauged five-dimensional supergravity. One can consistently set the  $\mathbf{1} + \mathbf{1} + \mathbf{10} + \overline{\mathbf{10}}$  of scalars to zero, at the same time as setting the  $\mathbf{6} + \overline{\mathbf{6}}$  of 2-form potentials to zero. The bosons that remain, namely gravity, the  $\mathbf{15}$  Yang-Mills fields, and the  $\mathbf{20'}$  of scalars, come just from the metric plus the self-dual 5-form sector of the original type IIB theory. We shall obtain complete results for the consistent embedding of this sub-sector of the gauged  $\mathcal{N} = 8$  theory, with all fifteen gauge fields  $A_{(1)}^{ij}$  of the  $SO(6)$ , and the twenty scalars that parameterize the full  $SL(6, \mathbb{R})/SO(6)$  sub-manifold of the complete scalar coset. These can be parameterized by a unimodular symmetric tensor  $T_{ij}$ .

Another way of expressing the truncation that we shall consider in this section is as follows. The type IIB theory itself can be consistently truncated in  $D = 10$  so that just gravity and the self-dual 5-form remain. The fifteen Yang-Mills gauge fields and twenty scalars that we retain in our ansatz are the full set of massless fields associated with Kaluza-Klein reduction from this ten-dimensional starting point. (The counting of massless fields is the same as the one that arises from a toroidal reduction from the same ten-dimensional starting point.) We shall see below that the self-duality condition on the 5-form plays an essential rôle in the consistency of the  $S^5$  reduction. This subsector of type IIB supergravity is particularly relevant for the AdS/CFT correspondence, because it is the metric and the self-dual 5-form that couple to the D3-brane as we will see in chapter IV.

We parameterize the fields for this truncated theory as follows. The twenty scalars, which are in the  $\mathbf{20'}$  representation of  $SO(6)$ , are represented by the symmetric unimodular tensor  $T_{ij}$ , where  $i$  is a 6 of  $SO(6)$ . The fifteen  $SO(6)$  Yang-Mills gauge

fields will be represented by the 1-form potentials  $A_{(1)}^{ij}$ , antisymmetric in  $i$  and  $j$ . The inverse of the scalar matrix  $T_{ij}$  is denoted by  $T_{ij}^{-1}$ . In terms of these quantities, we find that the Kaluza-Klein reduction ansatz is given by

$$d\hat{s}_{10}^2 = \Delta^{1/2} ds_5^2 + g^{-2} \Delta^{-1/2} T_{ij}^{-1} \mathcal{D}\mu^i \mathcal{D}\mu^j, \quad (3.13)$$

$$\hat{H}_{(5)} = \hat{G}_{(5)} + \hat{*}\hat{G}_{(5)}, \quad (3.14)$$

$$\begin{aligned} \hat{G}_{(5)} = & -g U \epsilon_{(5)} + g^{-1} (T_{ij}^{-1} * \mathcal{D} T_{jk}) \wedge (\mu^k \mathcal{D}\mu^i) \\ & - \frac{1}{2} g^{-2} T_{ik}^{-1} T_{j\ell}^{-1} * F_{(2)}^{ij} \wedge \mathcal{D}\mu^k \wedge \mathcal{D}\mu^\ell, \end{aligned} \quad (3.15)$$

$$\begin{aligned} \hat{*}\hat{G}_{(5)} = & \frac{1}{5!} \epsilon_{i_1 \dots i_6} \left[ g^{-4} U \Delta^{-2} \mathcal{D}\mu^{i_1} \wedge \dots \wedge \mathcal{D}\mu^{i_5} \mu^{i_6} \right. \\ & - 5g^{-4} \Delta^{-2} \mathcal{D}\mu^{i_1} \wedge \dots \wedge \mathcal{D}\mu^{i_4} \wedge \mathcal{D}T_{i_5 j} T_{i_6 k} \mu^j \mu^k \\ & \left. - 10g^{-3} \Delta^{-1} F_{(2)}^{i_1 i_2} \wedge \mathcal{D}\mu^{i_3} \wedge \mathcal{D}\mu^{i_4} \wedge \mathcal{D}\mu^{i_5} T_{i_6 j} \mu^j \right], \end{aligned} \quad (3.16)$$

where

$$\begin{aligned} U &\equiv 2T_{ij} T_{jk} \mu^i \mu^k - \Delta T_{ii}, & \Delta &\equiv T_{ij} \mu^i \mu^j, \\ F_{(2)}^{ij} &= dA_{(1)}^{ij} + g A_{(1)}^{ik} \wedge A_{(1)}^{kj}, & \mathcal{D}T_{ij} &\equiv dT_{ij} + g A_{(1)}^{ik} T_{kj} + g A_{(1)}^{jk} T_{ik}, \\ \mu^i \mu^i &= 1, & \mathcal{D}\mu^i &\equiv d\mu^i + g A_{(1)}^{ij} \mu^j, \end{aligned} \quad (3.17)$$

and  $\epsilon_{(5)}$  is the volume form on the five-dimensional space-time. Note that  $\hat{*}\hat{G}_{(5)}$  is derivable from the given expressions (3.13) and (3.15); we have presented it here because it is quite an involved computation. The coordinates  $\mu^i$ , subject to the constraint  $\mu^i \mu^i = 1$ , parameterize points in the internal 5-sphere. In obtaining the above ansatz we have been guided by previous results in the literature, including the  $S^4$  reduction ansatz from  $D = 11$  that was constructed in [20, 21].

It is consistent to truncate the fields of type IIB supergravity to the metric and self-dual 5-form  $\hat{H}_{(5)}$ . The ten-dimensional equations for motion for these fields can be read off from Eqs. (C.3)-(C.7). The ansatz presented above satisfies these equations

of motion if and only if the five-dimensional fields satisfy the equations

$$\begin{aligned}
\mathcal{D}(T_{ik}^{-1} * \mathcal{D}T_{kj}) &= -2g^2 (2T_{ik} T_{jk} - T_{ij} T_{kk}) \epsilon_{(5)} + T_{ik}^{-1} T_{\ell m}^{-1} * F_{(2)}^{\ell k} \wedge F_{(2)}^{mj} \\
&\quad - \frac{1}{6} \delta_{ij} \left[ -2g^2 (2T_{k\ell} T_{k\ell} - (T_{kk})^2) \epsilon_{(5)} + T_{pk}^{-1} T_{\ell m}^{-1} * F_{(2)}^{\ell k} \wedge F_{(2)}^{mp} \right], \\
\mathcal{D}(T_{ik}^{-1} T_{j\ell}^{-1} * F_{(2)}^{k\ell}) &= -2g T_{k[i}^{-1} * \mathcal{D}T_{j]k} - \frac{1}{8} \epsilon_{ijk_1 \dots k_4} F_{(2)}^{k_1 k_2} \wedge F_{(2)}^{k_3 k_4}, \tag{3.18}
\end{aligned}$$

together with the five-dimensional Einstein equation. These equations of motion can all be derived from the five-dimensional Lagrangian

$$\begin{aligned}
\mathcal{L}_5 &= R * \mathbb{1} - \frac{1}{4} T_{ij}^{-1} * \mathcal{D}T_{jk} \wedge T_{k\ell}^{-1} \mathcal{D}T_{\ell i} - \frac{1}{4} T_{ik}^{-1} T_{j\ell}^{-1} * F_{(2)}^{ij} \wedge F_{(2)}^{k\ell} - V * \mathbb{1} \\
&\quad - \frac{1}{48} \epsilon_{i_1 \dots i_6} \left( F_{(2)}^{i_1 i_2} F_{(2)}^{i_3 i_4} A_{(1)}^{i_5 i_6} - g F_{(2)}^{i_1 i_2} A_{(1)}^{i_3 i_4} A_{(1)}^{i_5 j} A_{(1)}^{j i_6} \right. \\
&\quad \left. + \frac{2}{5} g^2 A_{(1)}^{i_1 i_2} A_{(1)}^{i_3 j} A_{(1)}^{j i_4} A_{(1)}^{i_5 k} A_{(1)}^{k i_6} \right), \tag{3.19}
\end{aligned}$$

where the potential  $V$  is given by

$$V = \frac{1}{2} g^2 (2T_{ij} T_{ij} - (T_{ii})^2). \tag{3.20}$$

In (3.19) we have omitted the wedge symbols in the final topological term due to its cumbersome. The Lagrangian is in agreement with the one for five-dimensional gauged  $SO(6)$  supergravity in [80].

To see this, we look first at the ten-dimensional equation  $d\hat{H}_{(5)} = 0$ , with the ansatz given above. The terms involving structures of the form  $X_{(4)}^{ij} \wedge \mathcal{D}\mu^i \wedge \mathcal{D}\mu^j$ , where  $X_{(4)}^{ij}$  represents a 4-form in the five-dimensional space-time, give rise to the Yang-Mills equations above. Contributions of this kind come from  $d\hat{G}_{(5)}$  and also from the final term in  $d\hat{*}\hat{G}_{(5)}$ . The terms involving structures of the form  $X_{(5)}^{ij} \wedge (\mu^i \mathcal{D}\mu^j)$ , where  $X_{(5)}^{ij}$  represents a 5-form in the five-dimensional space-time, give rise to the scalar equations of motion in (3.18). Since  $\mu^i \mathcal{D}\mu^i = \mu^i d\mu^i = \frac{1}{2} d(\mu^i \mu^i) = 0$ , there is a trace subtraction in the scalar equation, and we read off  $X_{(5)}^{ij} - \frac{1}{6} \delta_{ij} X_{(5)}^{kk} = 0$  as the

five-dimensional equation of motion. Contributions of this structure come only from  $d\hat{G}_{(5)}$ . Finally, all other structures arising from calculating  $d\hat{H}_5 = 0$  vanish identically, without the use of any five-dimensional equations of motion. In deriving these results one needs to make extensive use of the Schoutens over-antisymmetrization identity,  $\epsilon_{[i_1 \dots i_6} V_{i_7]} = 0$ .

It is worth remarking that it is essential for the consistency of the reduction ansatz that the 5-form  $\hat{H}_{(5)}$  should be self-dual. One cannot simply consider a reduction ansatz for a ten-dimensional theory consisting of gravity plus a non-self-dual 5-form, whose ansatz is given by  $\hat{G}_{(5)}$  in (3.15). Although the Bianchi identity  $d\hat{G}_{(5)} = 0$  would give perfectly acceptable equations of motion for  $F_{(2)}^{ij}$  and  $T_{ij}$ , the field equation  $d\hat{*}\hat{G}_{(5)} = 0$  would produce the (unacceptable) constraint

$$\epsilon_{ijk_1 \dots k_4} F_{(2)}^{k_1 k_2} \wedge F_{(2)}^{k_3 k_4} = 0 . \quad (3.21)$$

It is only by combining  $\hat{G}_{(5)}$  and  $\hat{*}\hat{G}_{(5)}$  together into the self-dual field  $\hat{H}_{(5)}$  that a consistent five-dimensional result is obtained, with (3.21) now combining with terms from  $d\hat{G}_{(5)}$  to form part of the five-dimensional Yang-Mills equations given in (3.18). Notice that if one were to consider the reduction of a ten-dimensional theory with a non-self-dual 5-form there would be additional fields present in a complete massless truncation. These would comprise  $\mathbf{10} = \mathbf{4} + \mathbf{6}$  vector potentials and  $\mathbf{5} = \mathbf{1} + \mathbf{4}$  scalars. However the inclusion of these fields would still not achieve a consistent reduction ansatz, since the current (3.21), which is in the  $\mathbf{15}$  of  $SO(6)$ , could still not acquire an interpretation as a source term for the additional fields. Thus the additional requirement of self-duality seems to be essential for consistency. Of course, anti-self-duality would be equally good. It is interesting, therefore, that self-duality of the 5-form is apparently forced on us by the requirements of Kaluza-Klein consistency, if we try to “invent” a ten-dimensional theory that can be reduced on  $S^5$ . Thus once

again we see that supersymmetry and Kaluza-Klein consistency for sphere reductions seem to go hand in hand. Of course, the initial truncation of the type IIB theory to its gravity plus self-dual 5-form sector is itself a non-supersymmetric one, but the crucial point is that the consistency of the Kaluza-Klein  $S^5$  reduction is singling out a starting point that is itself a subsector of a supersymmetric theory.

The Kaluza-Klein  $S^5$  reduction that we have obtained here retains the full set of massless fields that can result from the reduction of gravity plus a self-dual 5-form in  $D = 10$ . In other words, after the initial truncation of the type IIB theory in  $D = 10$ , no further truncation of massless fields has been performed.

It should also be emphasized that it would be inconsistent to omit the fifteen  $SO(6)$  gauge fields when considering the embedding of the twenty scalars  $T_{ij}$ . This can be seen from the Yang-Mills equations in (3.18), which have a source term  $g T_{k[i}^{-1} * \mathcal{D} T_{j]k}$  appearing on the right-hand side. This is a quite different situation from a toroidal reduction, where it is always consistent to truncate to the scalar sector, setting the gauge fields to zero. The new feature here in the sphere reduction is that the scalar fields are charged under the gauge group. This is a general feature of all the sphere reductions, to  $D = 4$ ,  $D = 5$  and  $D = 7$ , and thus in all cases it is inconsistent to include the full set of scalar fields without including the gauge fields as well (see also [20, 21, 61]). (One can consistently truncate to the diagonal scalars in  $T_{ij}$ , setting all the gauge fields to zero, as in [32, 33], since then  $T_{k[i}^{-1} * \mathcal{D} T_{j]k} = 0$ .)

Our testing of the consistency of the reduction ansatz (3.13)-(3.16) has so far been restricted to checking the equations of motion for  $\hat{H}_{(5)}$ . A full testing of the consistency of the ten-dimensional Einstein equations would be quite involved, and will be addressed in future work. The experience in all previous work on consistent reductions is that although the actual checking of the higher-dimensional Einstein equations is the most difficult part from a computational point of view, the consistency

seems to be assured once it has been achieved for the equations of motion for the antisymmetric tensor fields, which itself is an extremely stringent requirement.

The  $S^5$  reduction ansatz that we have constructed here is also consistent if we include the dilaton  $\hat{\phi}$  and axion  $\hat{\chi}$  of the type IIB theory. These simply reduce according to the ansätze

$$\hat{\phi} = \phi , \quad \hat{\chi} = \chi , \quad (3.22)$$

where the unhatted quantities denote the fields in five dimensions. In this reduction they do not appear in the previous ansätze for the metric and self-dual 5-form, and in  $D = 5$  they just give rise to the additional  $SL(2, \mathbb{R})$ -invariant Lagrangian

$$\mathcal{L}_{(\phi, \chi)} = -\frac{1}{2} *d\phi \wedge d\phi - \frac{1}{2} e^{2\phi} *d\chi \wedge d\chi , \quad (3.23)$$

which is added to (3.19). Note in particular that  $\phi$  and  $\chi$  do not appear in the five-dimensional scalar potential  $V$ .

## CHAPTER IV

### SOME APPLICATIONS

#### A. Super Yang-Mills operators via AdS/CFT correspondence

There are two known ways to determine the SYM operators that correspond to given supergravity modes via AdS/CFT [16, 17, 18]. In the first approach [18, 81], one considers the representations of the super Lie algebra,  $SU(2, 2|4)$ , of the fields in SYM theory and IIB supergravity respectively, and matches the supergravity modes with the SYM operators by comparing the various quantum numbers. The other approach was proposed by Das and Trivedi [82]. They considered the lowest Kaluza-Klein mode of the NS-NS 2-form fields polarized along the D3-brane worldvolume. They worked out the corresponding SYM operators by expanding the Dirac-Born-Infeld (DBI) action plus Wess-Zumino (WZ) terms around an  $AdS_5 \times S^5$  background. They noted that the expansion around this background, as opposed to a flat background, is crucial in order to obtain the correct SYM operators. A similar method was used in [83, 84]

The relevance of curved backgrounds in AdS/CFT was already noticed in [85, 86] and was further motivated in [87]. Following [36], we consider another supergravity mode, and work out the SYM operators using the method presented in [82].

The ansatz given in Eqs. (3.13)-(3.16) is for a general  $S^5$  reduction, but for our discussion, we choose the space-time  $ds_5^2$  to be a five-dimensional anti-de Sitter space,  $AdS_5$ . Furthermore, to simplify our discussion, we consider configurations with

$A_{(1)}^{ij} = 0$ , which in turn enables us to put  $T_{ij}$  into diagonal form

$$T_{ij} = \text{diag}(X_1, X_2, X_3, X_4, X_5, X_6) ; \quad \prod_{i=1}^6 X_i = 1 . \quad (4.1)$$

We now adopt the parameterization used in [32]

$$X_i = \exp \left( -\frac{1}{2} \vec{b}_i \cdot \vec{\varphi} \right) , \quad (4.2)$$

where  $\vec{b}_i$  are the weight vectors of the fundamental representation of  $SL(6, \mathbb{R})$  which satisfy the following relations

$$\vec{b}_i \cdot \vec{b}_j = 8 \delta_{ij} - \frac{4}{3} , \quad \sum_{i=1}^6 \vec{b}_i = 0 , \quad \text{and} \quad \sum_{i=1}^6 (\vec{u} \cdot \vec{b}_i) \vec{b}_i = 8 \vec{u} , \quad (4.3)$$

where  $\vec{u}$  is an arbitrary vector and  $\vec{\varphi} = (\varphi_1, \varphi_2, \varphi_3, \varphi_4, \varphi_5)$  are five independent scalars.

The explicit expressions for  $\vec{b}_i$  can be taken as follows

$$\begin{aligned} \vec{b}_1 &= \left( 2, \frac{2}{\sqrt{3}}, \frac{2}{\sqrt{6}}, \frac{2}{\sqrt{10}}, \frac{2}{\sqrt{15}} \right) , \quad \vec{b}_2 = \left( -2, \frac{2}{\sqrt{3}}, \frac{2}{\sqrt{6}}, \frac{2}{\sqrt{10}}, \frac{2}{\sqrt{15}} \right) , \\ \vec{b}_3 &= \left( 0, -\frac{4}{\sqrt{3}}, \frac{2}{\sqrt{6}}, \frac{2}{\sqrt{10}}, \frac{2}{\sqrt{15}} \right) , \quad \vec{b}_4 = \left( 0, 0, -\sqrt{6}, \frac{2}{\sqrt{10}}, \frac{2}{\sqrt{15}} \right) , \\ \vec{b}_5 &= \left( 0, 0, 0, -\frac{8}{\sqrt{10}}, \frac{2}{\sqrt{15}} \right) , \quad \vec{b}_6 = \left( 0, 0, 0, 0, -\frac{10}{\sqrt{15}} \right) . \end{aligned} \quad (4.4)$$

With the diagonal form of  $T_{ij}$ , the Lagrangian (3.19) and equations (3.18) of motion up to fourth order in  $\vec{\varphi}$  are

$$\begin{aligned} e^{-1} \mathcal{L} &= -\frac{1}{2} (\partial \vec{\varphi}) \cdot (\partial \vec{\varphi}) + 12g^2 + \frac{1}{4} g^2 \sum_{i=1}^6 (\vec{b}_i \cdot \vec{\varphi})^2 + \frac{1}{24} g^2 \sum_{i=1}^6 (\vec{b}_i \cdot \vec{\varphi})^3 \\ &\quad - \frac{5}{192} g^2 \sum_{i=1}^6 (\vec{b}_i \cdot \vec{\varphi})^4 + \frac{1}{128} g^2 \left[ \sum_{i=1}^6 (\vec{b}_i \cdot \vec{\varphi})^2 \right]^2 \\ &= -\frac{1}{2} \partial_\mu \vec{\varphi} \cdot \partial^\mu \vec{\varphi} + 12g^2 + 2g^2 \vec{\varphi} \cdot \vec{\varphi} + V_3 + V_4 , \\ \square \vec{\varphi} &= -4g^2 \vec{\varphi} - \frac{1}{8} g^2 \sum_{i=1}^6 \vec{b}_i (\vec{b}_i \cdot \vec{\varphi})^2 + \frac{5}{48} g^2 \sum_{i=1}^6 \vec{b}_i (\vec{b}_i \cdot \vec{\varphi})^3 \end{aligned}$$



$$-\frac{1}{4}g^2\vec{\varphi}\sum_{i=1}^6(\vec{b}_i\cdot\vec{\varphi})^2, \quad (4.5)$$

where  $V_3$  and  $V_4$  are third-order and fourth-order polynomials in  $\vec{\varphi}$  respectively. We do not present here the explicit forms of  $V_3$  and  $V_4$  due to their cumbersome. However, their explicit forms can be found in [36].

In [37], two- and three-point correlators for various chiral primary operators were computed using a linear ansatz in [13]. As a consequence of using the linear ansatz, non-linear field redefinitions were required. The advantage of having a non-linear ansatz is obvious from (4.5): it renders such field redefinitions unnecessary. One can easily read off two- and three-point functions using the formulae in [88, 89].

Keeping only the terms linear in  $\varphi$ , the metric ansatz (3.13) can be written as

$$\begin{aligned} ds_{10}^2 \simeq & \left(1 - \frac{1}{4}\sum_i(\mu^i)^2\vec{b}_i\cdot\vec{\varphi}\right) ds_5^2 \\ & + \frac{1}{g^2}\left(1 + \frac{1}{4}\sum_i\vec{b}_i\cdot\vec{\varphi}(\mu^i)^2\right)\sum_i\left(1 + \frac{1}{2}\vec{b}_i\cdot\vec{\varphi}\right)(d\mu^i)^2. \end{aligned} \quad (4.6)$$

As previously mentioned, the ansatz quoted in Eq. (3.13) is for general  $S^5$  reductions. However, we choose the five-dimensional space-time to be  $AdS_5$  for our discussion. The structure of the D3-brane solution of type IIB supergravity in the near-horizon region is such that the “radius” of the  $AdS_5$  is the same as that of the internal sphere. Therefore we choose  $ds_5^2$  to be an  $AdS_5$  of radius  $1/g$ . Then we have

$$ds_5^2 + \frac{1}{g^2}\sum_i(d\mu^i)^2 = g^2r^2\sum_a(dx^a)^2 + \frac{1}{g^2r^2}\left(dr^2 + r^2\sum_i(d\mu^i)^2\right). \quad (4.7)$$

The metric ansatz (4.6) in its written form, i.e. in the  $\mu$ -coordinate system, does not make manifest the  $SO(6)$  covariance of the conformal field theory. A more suitable coordinate system is one that reveals the brane structure more transparently; not

surprisingly, such a coordinate system is of Cartesian type,

$$\mu^i = \frac{\Phi^i}{r} \quad \text{and} \quad r^2 = \sum_{i=1}^6 (\Phi^i)^2 . \quad (4.8)$$

Using the  $\Phi$ -coordinate system, one can rewrite (4.6) in the form

$$ds_{10}^2 = g^2 r^2 f \sum_a (dx^a)^2 + \frac{1}{g^2 r^2} \sum_{i,j=1}^6 g_{ij} d\Phi^i d\Phi^j , \quad (4.9)$$

where

$$\begin{aligned} f &\equiv 1 - \frac{1}{4r^2} \sum_i \vec{b}_i \cdot \vec{\varphi} (\Phi^i)^2 , \\ g_{ij} &\equiv \frac{1}{g^2 r^2} \left[ \delta_{ij} + \frac{1}{2} \vec{b}_i \cdot \vec{\varphi} \delta_{ij} - \frac{1}{r^2} \vec{b}_i \cdot \vec{\varphi} \Phi^i \Phi^j + \frac{1}{4r^2} \sum_k \vec{b}_k \cdot \vec{\varphi} (\Phi^k)^2 \delta_{ij} \right] . \end{aligned} \quad (4.10)$$

The action for D3 branes in a general type IIB background was studied in [90, 91, 92, 93]. For our discussion, it is enough to keep the metric and the 4-form field since the scalars come only from these. The relevant part of the action is

$$I = - \int d^4 \xi \sqrt{-\det(G_{ab} + F_{ab})} + \int \hat{C}_{(4)} , \quad (4.11)$$

where

$$\begin{aligned} G_{ab} &= \frac{\partial Z^M}{\partial \xi^a} \frac{\partial Z^N}{\partial \xi^b} E_M^A E_N^B \eta_{AB} , \\ C_{abcd} &= C_{MNPQ} \partial_a Z^M \partial_b Z^N \partial_c Z^P \partial_d Z^Q . \end{aligned} \quad (4.12)$$

It is the 4-form potential  $C_{(4)}$  that appears in (4.11), whereas the ansatz in Eqs. (3.15) and (3.16) was obtained in terms of the 5-form field strength,  $\hat{H}$ . It is possible to obtain  $C_{(4)}$  from Eqs. (3.15) and (3.16). Since we eventually consider only the leading order in the derivative expansion, it turns out that the contribution from  $C_{(4)}$  can be neglected. Therefore, we can concentrate solely on the contributions coming from the DBI action.

To obtain  $G_{ij}$ , we substitute (4.9), (4.10) and the super vielbeins obtained in [94, 95] into the first equation of (4.12). The result is

$$G_{ab} = g^2 r^2 f \eta_{ab} + g_{ij} \partial_a \Phi^i \partial_b \Phi^j + F_{ab} . \quad (4.13)$$

There are no fermionic terms in the above formula. The reason is that fermionic fields do not have any contributions with conformal dimension 2, which is a relevant dimension to our discussion.

After some algebra, the relevant part of the action is given by

$$\begin{aligned} I[\phi] = & - \int -\frac{g^4 r^2}{2} \sum_i \vec{b}_i \cdot \vec{\varphi} (\Phi^i)^2 + \frac{1}{2} \sum_{ij} \left( \frac{1}{2} \vec{b}_i \cdot \vec{\varphi} \delta_{ij} - \frac{1}{r^2} \vec{b}_i \cdot \vec{\varphi} \Phi^i \Phi^j \right) \partial_a \Phi^i \partial^a \Phi^j \\ & + \dots , \end{aligned} \quad (4.14)$$

where the ellipses refer to the terms with higher numbers of derivatives.

The discussion of the full 20 scalars  $T_{ij}$ , without imposing  $A_{(1)}^{ij} = 0$ , goes very similarly, since eventually we will be interested only in terms that are coupled to  $T_{ij}$ , but do not have any factors of  $A_{(1)}^{ij}$ . In fact the computation is almost identical, except that one needs the new parameterization

$$T_{ij} \equiv (e^S)_{ij} , \quad (4.15)$$

where  $S_{ij}$  is symmetric and traceless. Using this parameterization, one gets

$$\begin{aligned} I[S_{ij}] = & - \int g^4 r^2 \Phi^i \Phi^j S_{ij} + \frac{1}{2} \left( -\partial_a \Phi^i \partial^a \Phi^j + \frac{1}{r^2} \left[ \Phi^i \Phi^k \partial_a \Phi^j \partial^a \Phi^k + i \leftrightarrow j \right] \right) S_{ij} \\ & - (\text{trace}) . \end{aligned} \quad (4.16)$$

To leading order in the derivative expansion, we have

$$I[S_{ij}] = - \int g^4 r^2 \left( \Phi^i \Phi^j - \frac{1}{6} \delta^{ij} \Phi^k \Phi_k \right) S_{ij} + \dots . \quad (4.17)$$

As one can easily show,  $S_{ij} \sim \frac{1}{r^2}$  in the boundary region, i.e.  $r \rightarrow \infty$ . Therefore we impose the following boundary condition,

$$S_{ij} \equiv \frac{1}{r^2} S_{ij}^o, \quad (4.18)$$

which leads to the correct CFT operator.

$$\mathcal{O}^{ij} \equiv \left( \Phi^i \Phi^j - \frac{1}{6} \delta^{ij} \Phi^k \Phi_k \right) + \dots. \quad (4.19)$$

The result is given in Eq. (4.19) is in agreement with the CFT operators obtained based on the conformal symmetry argument [18].

#### B. Supergravity duals of $\mathcal{N} = 2$ SCFTs in $D = 3$ and $D = 5$

We start this section by reviewing the general idea of supergravity description of field theories on a curved manifold presented in [39]. Generally, if we put a supersymmetric field theory on a curved manifold  $\mathcal{M}$ , we will break supersymmetry because we will not have a covariantly constant spinor which obeys  $(\partial_\mu + \omega_\mu) \epsilon = 0$ . If the supersymmetric field theory has a global R-symmetry, we then can couple the theory to an external gauge field that coupled to R-symmetry current and the spinor obeys the equation  $(\partial_\mu + \omega_\mu - A_\mu) \epsilon = 0$ . Obviously, if we choose the external gauge field to be equal to the spin connection  $\omega_\mu = A_\mu$ , then we can find a covariantly constant spinor. The resulting theory is the so-called “twisted” theory, since we can view the coupling to the external gauge field as effectively changing the spins of all fields. The supersymmetry parameter becomes a scalar. This is precisely the way that branes wrapping on non-trivial cycles in M-theory or string compactifications manage to preserve some supersymmetries [38]. A supersymmetric cycle is defined by the condition that a worldvolume theory is supersymmetric [96, 97]. The condition to have

a supersymmetric (p+1)-cycle is

$$\left(1 - \frac{i}{(p+1)!} h^{-1/2} \varepsilon^{\alpha_1 \dots \alpha_{p+1}} \partial_{\alpha_1} X^{m_1} \dots \partial_{\alpha_{p+1}} X^{m_{p+1}} \Gamma_{m_1 \dots m_{p+1}}\right) \Psi = 0, \quad (4.20)$$

where  $\Psi$  is a covariantly constant ten-dimensional spinor,  $h_{\alpha\beta}$  is an induced metric on p-brane and  $\Gamma_{m_1 \dots m_{p+1}}$  are ten-dimensional  $\Gamma$ -matrices. The cycles satisfying Eq. (4.20) minimize the p-brane worldvolume [97]. In the case of branes wrapping on cycles,  $\mathcal{M}$  is the worldvolume geometry of the cycle and the external gauge field takes into account the fact that the normal to the cycle forms a non-trivial bundle, the normal bundle, and  $A_\mu$  is the connection on this normal bundle. The condition that the cycle preserves some supersymmetries then boils down to the condition that the spin connection is equal to the gauge connection, which is exactly the equation mentioned earlier. So if we are interested in understanding field theories arising on the branes wrapping on some non-trivial cycles, we will then have to study these twisted field theories.

In this section we construct various supergravity solutions, some of which are duals to  $\mathcal{N} = 2$  SCFTs in three and five dimensions. These cases provide new examples of AdS/CFT duality involving twisted field theories.

### 1. Supergravity duals of three-dimensional $\mathcal{N} = 2$ SCFT

In this subsection, we focus on the D4-D8 system, which was studied from the gauge theory side in [98] and from the gravity/CFT viewpoint in [99, 100]. We will wrap D4-brane of the D4-D8 system on 2- and 3-cycles. These systems are the gravity duals of superconformal field theories in five dimensions with 8 supercharges [98], which in the infrared (IR) flow to superconformal field theories in three dimensions with 4 supercharges, which is the  $\mathcal{N} = 2$  theory in three dimensions [101, 102, 103, 104, 105].

We start with a brief review of D4-D8 system [100]. Consider type I theory on

$R^9 \times S^1$  with  $N$  coinciding D5-branes wrapping on the circle. The six-dimensional field theory on the worldvolume of D5-branes is  $\mathcal{N} = 1$  with  $Sp(N)$  gauge group [106]. This theory has one hypermultiplet in the antisymmetric representation of  $Sp(N)$  from Dirichlet-Dirichlet sector and 16 hypermultiplets in the fundamental representation from the Neuman-Dirichlet sector. T-dualizing D5-branes on the circle results in type I' theory compactified on the interval  $S^1/\mathbb{Z}_2$  with two orientifolds (O8-planes) located at fixed points [107]. The D5-branes become D4-branes and there are 16 D8-branes located at points on the interval. The existence of 16 D8-branes is necessary in order for R-R charge to be conserved and furthermore the theory is anomaly-free [108, 109]. The relative positions of the D8-branes with respect to the D4-brane correspond to masses for the hypermultiplet in the fundamental representation. The hypermultiplet in the antisymmetric representation is massless.

The field theory on the worldvolume of D4-brane is a five-dimensional supersymmetric field theory [110, 111]. The Lorentz group of D4-brane worldvolume is  $SO(4, 1)$ . The spinor representation of  $SO(1, 4)$  is pseudoreal and is four-dimensional. This theory in five dimensions is related by dimensional reduction to the  $\mathcal{N} = 1$  theory in six dimensions described above. All these theories have an  $SU(2)_R$  symmetry of automorphism algebra which can be associated with the gauged symmetry in the gravitational set-up of [26]. The vector multiplet in five dimensions consists of a real scalar component, a vector field and a spinor, while the hypermultiplet comprises four real scalars and a spinor. This theory has a Coulomb branch when the real scalar has a vacuum expectation value (VEV), and a Higgs branch when the scalar in the hypermultiplet is excited. The theory on the D4-brane has  $Sp(N)$  gauge group with one vector multiplet (whose scalar component describes the Coulomb branch  $\mathbb{R}^+$ ) and hypermultiplets whose first components describe the Higgs branch. The global symmetries of the theory are  $SU(2)_R \times SU(2) \times SO(2N_f) \times U(1)$ . The  $SU(2)_R$  fac-

tor is the R-symmetry (the supercharges and the scalars in the hypermultiplets are doublets under this group). The other  $SU(2)$  is associated with the hypermultiplet in the antisymmetric representation which is only present if there are more than one D4-brane (e.g.  $N > 1$ ). The  $SO(2N_f)$  is the symmetry group of the  $N_f$  massless hypermultiplets in the fundamental representation, and the  $U(1)$  is associated with the instanton number. When the D4-brane is in the origin of the Coulomb branch we have a fixed point and the global symmetry is enhanced to  $SU(2) \times E_{N_f+1}$ . The gravitational system is given by  $N_f$  D8-branes situated on a O8-plane,  $(16 - N_f)$  D8-branes in another fixed plane, and a D4-brane which can move between them. The position of the D4-brane is parameterized by the scalar in the vector multiplet. If  $\langle \phi \rangle$  is not zero, we have a theory with a  $U(1)$  symmetry with  $N_f$  “electrons”, on the fixed planes; the theory recovers its  $SU(2)$  R-symmetry, and it will have  $N_f$  quarks.

The gravity theory is a fibration of  $AdS_6$  over  $S^4$ , with isometries  $SO(2,5) \times SU(2) \times SU(2)$ . The six-dimensional  $SU(2)$ -gauged supergravity has an  $AdS_6$  vacuum solution. It can be up-lifted to massive IIA theory [22] leading to the configuration described above [48]. Other solutions in massive type IIA supergravity were obtained in [112, 113]. Here we will consider solutions of Romans’ theory which, when uplifted, will give ten-dimensional configurations with  $N_f = 0$ . These theories were studied in depth in [114] in the context of algebraic geometry. There, the authors studied the moduli spaces of the type I’ string and the heterotic string. It would be interesting to understand the case of  $N_f \neq 0$  and see whether there is a gravity solution dual to the theory with enhanced symmetry  $E'_1$ . It would be also useful to clarify the role of D0-branes in our gravity solutions. In principle they should correspond to excitations of the six-dimensional Abelian gauge field.

In order to analyze the flow of the theory from five dimensions to a supercon-

formal theory in three dimensions with 4 supercharges, we consider a configuration where the geometry “loses” two dimensions at low energies. The idea is to start with a six-dimensional gauged supergravity theory which has an  $AdS_6$  vacuum with 8 preserved supercharges [50]. This vacuum solution is dual to the five-dimensional SCFT mentioned above and its ten-dimensional interpretation is given by a D4-D8 system. When one moves to the IR of the gauge theory (flowing in the radial coordinate on the gravity dual) two of the dimensions of the theory become very small and no low-energy massless modes are excited on this two-dimensional space. Therefore, the gauge theory is effectively three-dimensional. Further, since the D4-brane is wrapped on a curved surface, we need to twist the theory. The effect of this twisting is the breaking of some supersymmetries, as can be seen from the projections of the Killing vectors.

Under the group  $SO(1, 4) \times SO(3)_R$  the fields in the vector multiplet on the D4-brane worldvolume transform as  $(\mathbf{1}, \mathbf{1})$  for the scalar field,  $(\mathbf{3}, \mathbf{1})$  for the gauge field and  $(\mathbf{4}, \mathbf{2})$  for the fermion. In the IR, the metric is taken to have  $SO(1, 2) \times SO(2)_D$  symmetry. The twisting involves “mixing” the  $SO(2)$  group of the metric with an  $SO(2)$  subgroup of  $SO(3)_R$ . This mixing is responsible for the breaking of some supersymmetries because the only spinors that can be defined on the curved manifold are those that have scalar properties on the curved part. In fact, it breaks 1/2 of the supersymmetries of the five-dimensional SCFT.

Below we construct a gravity dual to the five-dimensional SCFT theory by finding a solution of Romans’ theory with excited  $F_{\mu\nu}^I$  fields. Then, we find a fixed-point solution for the massive field configuration. We postpone the study of the massless case to the next subsection.



Let us consider the metric ansätze of the form

$$ds^2 = e^{2f} (dt^2 - dr^2 - dz^2 - dv^2) - \frac{e^{2h}}{y^2} (dx^2 + dy^2) , \quad (4.21)$$

for  $AdS_4 \times H_2$  and

$$ds^2 = e^{2f} (dt^2 - dr^2 - dz^2 - dv^2) - e^{2h} (d\theta^2 + \sin^2 \theta d\varphi^2) , \quad (4.22)$$

for  $AdS_4 \times S^2$ . The only non-vanishing non-Abelian gauge potential components are taken to be

$$A_x^{(3)} = \frac{a}{y} \quad \text{for } AdS_4 \times H_2 \text{ case} , \quad (4.23)$$

$$A_\varphi^{(3)} = -a \cos \theta \quad \text{for } AdS_4 \times S^2 \text{ case} . \quad (4.24)$$

One can treat both cases together by introducing a parameter  $\lambda$  that can take two values

$$\lambda = +1 \rightarrow H_2 , \quad \lambda = -1 \rightarrow S^2 . \quad (4.25)$$

We impose the following projections

$$\Gamma_{45}(T^{(3)})_i{}^j \epsilon_j = \frac{\lambda}{2} \epsilon_i , \quad \Gamma_2 \Gamma_7 \epsilon_i = \epsilon_i . \quad (4.26)$$

A solution must satisfy the equations obtained by setting to zero the supersymmetry transformations for gauginos and gravitinos. After some algebra, they lead to

$$\varphi' = \frac{1}{4\sqrt{2}} e^f [g e^\varphi - 3m e^{-3\varphi} + 2a \lambda e^{-2h-\varphi}] , \quad (4.27)$$

$$h' = \frac{1}{4\sqrt{2}} e^f [-g e^\varphi - m e^{-3\varphi} + 6a \lambda e^{-2h-\varphi}] , \quad (4.28)$$

$$f' = -\frac{1}{4\sqrt{2}} e^f [g e^\varphi + m e^{-3\varphi} + 2a \lambda e^{-2h-\varphi}] , \quad (4.29)$$

where  $\varphi \equiv \phi/\sqrt{2}$ .

A fixed-point solution of these equations is given by

$$\begin{aligned} e^{4\varphi} &= \frac{2m}{g} , \\ e^{-2h} &= \sqrt{\frac{gm}{8}} \frac{1}{a\lambda} , \\ e^{-f(r)} &= \frac{g e^{\varphi}}{2\sqrt{2}} r . \end{aligned} \tag{4.30}$$

In fact, it only happens if  $\lambda a > 0$ . This solution satisfies the second order equations derived from the six-dimensional Lagrangian.

We can analyze the structure of the solutions of the system above near the UV of the theory ( $r = 0$ ). Indeed, we can see that an expansion leads to the following behavior for the different fields

$$f \approx -\log(r) + c_1 r^2 + \dots, \quad g \approx -\log(r) + c_2 r^2 + \dots, \quad \phi \approx c_3 r^2 + \dots, \tag{4.31}$$

where  $c_i$ s are constants. The interpretation of these expansions is that the scalar field  $\phi$  describes the coupling of the five-dimensional field theory to an operator of conformal weight  $\Delta = 3$  and mass squared  $m^2 = -6$  [115]. The operator is not the highest component of its supermultiplet, thus it is a deformation that breaks some supersymmetry.

The ten-dimensional metric corresponding to the fixed-point solution described above is easily obtained by using the results of [22],

$$\begin{aligned} ds_{10}^2 &= (3mg^2)^{-1/8} (\sin \chi)^{1/12} X^{1/8} \left\{ \frac{\Delta^{3/8} e^{2f_0}}{2r^2} (-dt^2 + dr^2 + dv^2 + dz^2) \right. \\ &\quad + \frac{\sqrt{2}\Delta^{3/8}}{\sqrt{g^3 m} y^2} (dx^2 + dy^2) + \Delta^{3/8} \sqrt{2mg^3} d\chi^2 \\ &\quad \left. + \frac{\Delta^{-5/8}}{gX} \cos^2 \chi \left[ \left( \sigma_1 - \frac{dx}{y} \right)^2 + \sigma_2^2 + \sigma_3^2 \right] \right\} . \end{aligned} \tag{4.32}$$

The metric has symmetries of the form  $SO(2) \times SO(3)$ . The corresponding expression

for the R-R and NS-NS fields and the ten-dimensional dilaton can be read off from (E.22).

Although we could not find an exact solution, we can carry out an analysis similar to [47]. Actually, the two cases  $\lambda = \pm 1$  can be gathered again. Thus defining  $F \equiv u^2 e^{-2\varphi}$ ,  $u \equiv e^{2h(r)}$  and using Eqs. (4.27)-(4.28) we obtain the following differential equation

$$\frac{dF}{du} = \left( \frac{3gu^4 - mF^2 - 10a\lambda uF}{gu^4 + mF^2 - 6a\lambda uF} \right) \frac{F}{u}. \quad (4.33)$$

For the case  $\lambda = +1$  (hyperbolic plane), we can solve this equation in the approximations when  $u \rightarrow \infty$  and  $u \rightarrow 0$ . In fact, in the UV one has

$$F \approx u^2 + \dots, \quad e^{-2\varphi} \approx 1 + \dots, \quad (4.34)$$

and the metric results to be the  $AdS_6$  limit of our original metric. In the IR we have

$$F \approx F_0 u^{5/3} + \dots, \quad e^{-2\varphi} \approx F_0 u^{-1/3} + \dots, \quad (4.35)$$

and the metric is

$$ds_6^2 = \frac{1}{u^{1/3}}(dt^2 - dz^2 - dv^2) - \frac{u}{y^2}(dx^2 + dy^2) - \frac{2}{9F_0 a u^{1/3}} du^2. \quad (4.36)$$

When up-lifted, the singularities turn out to be of the “bad” type. They correspond to the curves flowing to the origin of the plots in Fig. 1. On the other hand, in both cases of the hyperbolic plane and sphere there are other kinds of “bad” singularities that correspond to the orbits going like  $F \approx \frac{1}{u}$  as  $u$  approaches to zero. Particularly, one can see this kind of behavior in Fig. 2. For the two-sphere we have the set of orbits depicted in Fig. 2.

This also constitutes a part of the IR behavior of the flow. As we can see in Fig. 1 and Fig. 2, there is a smooth solution interpolating between the UV and the




Fig. 1. Orbits for the case  $AdS_4 \times H_2$ . In the UV limit  $F$  behaves as  $u^2$  leading to an  $AdS_6$ -type region. On the other hand, it flows to “bad” singularities (BS) in the IR. We have used  $g = 3m$  with  $m = \sqrt{2}$ , and  $a = 1$ .

IR limits of the system, thus realizing our initial set up. The fixed-point solution is out of the range of this plot.

In summary, we have found a fixed-point solution for the hyperbolic plane. In the UV, both geometries flow to their corresponding asymptotic  $AdS_6$ . In the IR limit all the singularities in both geometries are of the “bad” type according to the criterion of [39]. Similar behavior is expected for the sphere. Note that the solutions studied here only represent a special case of solutions that, in general, have the form of a Laurent series. The fact that these solutions have non-acceptable singularities can be easily seen from the construction of an effective potential corresponding to this effective four-dimensional system, and following the analysis outlined by Gubser in [116] one can see that these kinds of solutions will not be interpretable as the IR of a gauge theory. However, this does not exclude the possibility of finding a different solution that could be interpreted as the transcendental solution found in [47].




Fig. 2. We show the behavior of the orbits for the case  $S^2$ . Similar comments as in the previous case are also valid here. For small  $u$ -values we can see the region corresponding to  $F \approx \frac{1}{u}$ .

## 2. A non-Abelian solution in massive type IIA theory

In this subsection we present a non-Abelian solution in the massive IIA theory. We begin with a gravity dual of a five-dimensional SCFT with 8 supercharges that flows through the dimensions to a two-dimensional (1,1) CFT with 2 supercharges at the IR. The field content consists of a real scalar, a gauge field and a corresponding fermionic partner. The idea is to find a compactification from six-dimensional supergravity on  $AdS_3 \times H_3$ . Under  $SO(1,4) \times SO(3)_R$  the charges transform as  $(\mathbf{4}, \mathbf{2})$  and  $(\bar{\mathbf{4}}, \mathbf{2})$ . After the twisting  $SO(1,4) \times SO(3)_R \rightarrow SO(1,1) \times SO(3) \times SO(3)_R \rightarrow SO(1,1) \times SO(3)_D$  we are left with 2 supercharges which are scalars under the diagonal subgroup.  $SO(3)_D$  results from the identification of the normal bundle of the D4-brane world-volume with the spin bundle of  $\Sigma_3$ . They are the two “twisted” supercharges that survive the twisting process. This solution is similar to the solution obtained in [47] for massive seven-dimensional gauged supergravity. In that case the solution up-lifts to M-theory. This suggests some connection between theories in the line of [117] that merits further exploration.

Our configuration is

$$ds^2 = e^{2f}(dt^2 - dr^2 - du^2) - \frac{e^{2h}}{y^2}(dx^2 + dz^2 + dy^2), \quad A_x^{(1)} = \frac{a}{y}, \quad A_z^{(3)} = \frac{b}{y}. \quad (4.37)$$

Similarly, an  $AdS_3 \times S^3$  for the case  $\lambda = -1$  can also be considered. The projections (all the indices are flat indices) are given by

$$\Gamma_{65}(T^{(3)})_i{}^j \epsilon_j = \frac{1}{2} \epsilon_i, \quad \Gamma_{64}(T^{(2)})_i{}^j \epsilon_j = \frac{1}{2} \epsilon_i, \quad \Gamma_{45}(T^{(1)})_i{}^j \epsilon_j = \frac{1}{2} \epsilon_i, \quad (4.38)$$

and

$$\Gamma_7 \Gamma_2 \epsilon_i = \epsilon_i, \quad (4.39)$$

we use  $\xi = e^{\phi/\sqrt{2}}$ . The BPS equations are

$$h' = -\frac{1}{4\sqrt{2}} e^f [g\xi + m\xi^{-3} - 10\lambda a\xi^{-1}e^{-2h}], \quad (4.40)$$

$$\phi' = \frac{1}{4\sqrt{2}} e^f [g\xi - 3m\xi^{-3} + 6\lambda a\xi^{-1}e^{-2h}], \quad (4.41)$$

$$f' = -\frac{1}{4\sqrt{2}} e^f [g\xi + m\xi^{-3} + 6\lambda a\xi^{-1}e^{-2h}], \quad (4.42)$$

and  $ab = 1/g$ . A fixed-point solution is

$$e^{-2h} = \frac{\sqrt{m}g^3}{2\sqrt{6}}, \quad \xi = \left(\frac{3m}{2g}\right)^{1/4}. \quad (4.43)$$

This configuration can be uplifted to massive type IIA in ten dimensions using the results in [22]

$$\begin{aligned} ds_{10}^2 = & \left(\frac{1}{3mg^2}\right)^{1/8} (\sin\chi)^{1/12} X^{1/8} \left\{ \frac{\Delta^{3/8}}{2r^2}(-dt^2 + dr^2 + dv^2) \right. \\ & + \frac{e^{2h}}{2y^2} \Delta^{3/8} (dx^2 + dz^2 + dy^2) + g^2 \xi \Delta^{3/8} d\chi^2 \\ & \left. + \frac{\Delta^{-5/8}}{gX} \cos^2\chi \left[ \left(\sigma_1 - \frac{dx}{y}\right)^2 + \sigma_2^2 + \left(\sigma_3 - \frac{dz}{y}\right)^2 \right] \right\}, \quad (4.44) \end{aligned}$$

where  $\sigma_i$  are the three left-invariant forms in the 3-sphere whose explicit forms are

given in Eq. (3.5). We can see that the metric has an  $SO(2) \times SO(3)$  invariance.

Since we are not able to integrate the above BPS system, we will repeat the analysis of the singularities as we did in the previous section. Again, both cases (hyperbolic plane and sphere) are treated together by using the parameter  $\lambda = \pm 1$ . Proceeding as above we get the following first order differential equation

$$\frac{dF}{du} = \left( \frac{3gu^4 - mF^2 - 14a\lambda uF}{gu^4 + mF^2 - 10a\lambda uF} \right) \frac{F}{u} . \quad (4.45)$$

When  $\lambda = +1$  we solve this equation in two approximations,  $u \rightarrow \infty$  and  $u \rightarrow 0$ . The large- $u$  approximation leads to the corresponding asymptotic  $AdS_6$  whereas the other case gives

$$F \approx F_0 \frac{1}{u} + \dots , \quad e^{-2\varphi} \approx \frac{F_0}{u^3} + \dots , \quad (4.46)$$

$$F \approx -2\frac{a}{m}u + \dots , \quad e^{-2\varphi} \approx -\frac{2a}{mu} + \dots \quad (4.47)$$

$$F \approx F_0 u^{7/5} + \dots , \quad e^{-2\varphi} \approx F_0 u^{3/5} + \dots . \quad (4.48)$$

Note that all the singularities are of the “bad” type in the IR. They correspond to the curves flowing to the origin of the plots in Fig. 3. Here we should make a comment similar to that in the previous section: Gubser’s criteria tells us to construct the effective potential of the three-dimensional gravity theory. With this potential we can see that the solutions in the form of a Laurent series shown above will not have an interpretation as the IR limit of a gauge theory.

We can study a problem similar to the one in the previous section. One can write down a similar ansatz for the metric, i.e. a wrapped product of  $AdS_4 \times \Sigma_2$ , where  $\Sigma_2$  is a hyperbolic plane or sphere, as before. The system of Eqs. (4.27)-(4.29) can be studied in the massless case, i.e.  $m = 0$ , while  $g$  and  $a$  are non-vanishing quantities.




Fig. 3. Behavior of the orbits for the non-Abelian massive case. The  $AdS_6$ -type region is at the UV, i.e. for large  $u$  and  $F$ -values. In the IR it flows to “bad” singularities (BS). The cross indicates the fixed point-solution aforementioned. We again set  $g = 3m$ ,  $m = \sqrt{2}$ , and  $a = b = \lambda = 1$ .

We have the solution

$$f = -\varphi, \quad \varphi = \frac{g}{4\sqrt{2}}r + \frac{1}{8}\log(r), \quad h = -\frac{g}{4\sqrt{2}}r + \frac{3}{8}\log(r). \quad (4.49)$$

Therefore, the six-dimensional metric is given by

$$ds^2 = \frac{1}{r^{1/4}} e^{-\frac{gr}{2\sqrt{2}}} [dt^2 - dr^2 - dz^2 - dv^2 - r d\Omega_\lambda^2]. \quad (4.50)$$

This metric has a singularity. One way to resolve it is to look for a solution that has more degrees of freedom excited such as a non-Abelian solution. Another way to have an acceptable gravity solution is to construct a black hole solution such that the singularity is hidden by the horizon [118, 119, 120]. In the next section we will discuss a resolution of the singularity in the  $S^2$  case, i.e.  $\lambda = -1$  in Eqs. (4.27)-(4.29).

### 3. A non-Abelian solution

We would like to present a non-Abelian solution that resolves the “bad” singularity of the solution in Eq. (4.50). This solution is useful for the study of the gravity dual for



the three-dimensional  $\mathcal{N} = 2$  super Yang-Mills theory. This solution, which relates to the solution in [40], represents a smeared NS5-brane on  $S^2$  after a T-duality. In the IR, the theory living on the brane will be an  $\mathcal{N} = 2$  super Yang-Mills theory in three dimensions plus Kaluza-Klein modes that do not decouple. The solution in the string frame is

$$\begin{aligned}
ds^2 &= 2 [dt^2 - dr^2 - d\vec{x}_2^2 - e^{2h}(d\theta^2 + \sin^2 \theta d\varphi^2)] , \\
F &= -w' \sigma^2 dr \wedge d\theta + w' \sin \theta \sigma^1 dr \wedge d\varphi + (w^2 - 1) \sigma^3 \sin \theta d\theta \wedge d\varphi , \\
\phi_S &= \frac{1}{2} \log \left[ \frac{\sinh(r)}{R(r)} \right] , \quad w(r) = \frac{r}{\sinh(r)}, w' = \frac{dw}{dr} , \\
e^{2h} &= R(r)^2, R(r) = \sqrt{2r \coth(r) - \left[ \frac{r}{\sinh(r)} \right]^2 - 1} .
\end{aligned} \tag{4.51}$$

This is the same solution obtained by Chamseddine and Volkov in [121, 122].

The relations between the scalars and metric in the Einstein and string frames are

$$\varphi = -\frac{1}{2} \phi_S , \quad g_{\mu\nu}(E) = e^{\phi_S} g_{\mu\nu}(S) .$$

In Einstein frame the solution is

$$ds_E^2 = 2e^{\phi_S} (-dt^2 + dr^2 + d\vec{x}_2^2) + e^{2G(r)} (d\theta^2 + \sin^2 \theta d\varphi^2) , \tag{4.52}$$

where the dilaton is given by

$$\begin{aligned}
\phi_S &= \frac{1}{2} \log \left[ \frac{\sinh(r)}{R(r)} \right] , \\
R(r) &= \sqrt{2r \coth(r) - w(r)^2 - 1} , \quad e^{2G} = R(r)^2 , \quad w(r) = \frac{r}{\sinh(r)} .
\end{aligned} \tag{4.53}$$

The field strength components are

$$\begin{aligned}
F_{(2)}^1 &= w' \sin \theta dr \wedge d\varphi , \\
F_{(2)}^2 &= -w' dr \wedge d\theta ,
\end{aligned}$$

$$F_{(2)}^3 = (w^2 - 1) \sin \theta d\theta \wedge d\varphi . \quad (4.54)$$

Collecting all, we have the following ten-dimensional solution:

$$\begin{aligned} d\hat{s}_{10}^2 &= e^{-\frac{3\sqrt{2}}{8}\phi} \left[ -dt^2 + dr^2 + d\vec{x}_2^2 + e^{2G}(d\theta^2 + \sin^2 \theta d\varphi^2) + \frac{1}{g^2} \sum_{i=1}^3 \left( \sigma^i - \frac{g}{\sqrt{2}} A_{(1)}^i \right)^2 \right] \\ &\quad + e^{\frac{5\sqrt{2}}{8}\phi} dZ^2 , \\ \hat{F}_{(4)} &= \left( -\frac{1}{g^2} h^1 \wedge h^2 \wedge h^3 + \frac{1}{\sqrt{2}g} F_{(2)}^i \wedge h^i \right) \wedge dZ , \\ \hat{\phi} &= \frac{1}{\sqrt{2}} \phi = -\frac{1}{4} \log \left[ \frac{\sinh(r)}{R(r)} \right] , \end{aligned} \quad (4.55)$$

where  $h^i = \sigma^i - \frac{g}{\sqrt{2}} A_{(2)}^i$  and  $\sigma^i$  are left-invariant 1-forms in the  $S^3$  as given in Eqs. (3.4)-(3.5).

#### 4. A black hole solution in the massless theory

The solutions of the massless six-dimensional Romans' theory, the string frame metric, and 2-form ansätze are

$$\begin{aligned} ds^2 &= \frac{f(r)}{r} dt^2 - \frac{1}{r f(r)} dr^2 - R^2 (d\theta_1^2 + \sin^2 \theta_1 d\varphi_1^2) - dx^2 - dy^2, \\ F_2 &= h(r) dr \wedge dt + \frac{R Q_m}{\sqrt{2}} \sin \theta_1 d\theta_1 \wedge d\varphi_1 , \end{aligned} \quad (4.56)$$

where  $f$  and  $h$  are functions of  $r$  while  $\gamma$  and  $Q_m$  are constants.

The solution in the string frame is

$$h(r) = \frac{Q_e}{r^2} , \quad f(r) = -M + \frac{2Q_e^2}{r} + \left( \frac{g^2}{2} + \frac{Q_m^2}{R^2} \right) r , \quad \phi(r) = \frac{1}{2} \log(r) . \quad (4.57)$$

Similar solutions were discussed previously in [62]. Our solution has a different geometry and carries both electric and magnetic charges.

One can up-lift the above solution to type IIA theory using the ansätze (2.67),

and the results are

$$\begin{aligned}
d\hat{s}_{10}^2 &= -\frac{1}{2} r^{-1/8} \left[ \frac{f(r)}{\sqrt{r}} dt^2 - \frac{1}{\sqrt{r} f(r)} dr^2 - R^2 \sqrt{r} (d\theta_1^2 + \sin^2 \theta_1 d\varphi_1^2) \right] \\
&\quad + \frac{1}{2} r^{3/8} (dx^2 + dy^2) + r^{-5/8} dZ^2 \\
&\quad + \frac{r^{3/8}}{g^2} \left[ (\sigma^1)^2 + (\sigma^2)^2 + \left( \sigma^3 + \frac{g Q_e}{\sqrt{2} r} dt + \frac{g R Q_m}{2} \cos \theta_1 d\varphi_1 \right)^2 \right], \\
\hat{\phi} &= -\frac{1}{4} \log(r), \\
\hat{F}_4 &= -\frac{1}{g^2} \sigma^1 \wedge \sigma^2 \wedge h^3 \wedge dZ + \frac{1}{\sqrt{2} g} F_2 \wedge h^3 \wedge dZ, \\
h^3 &= \sigma^3 + \frac{g Q_e}{\sqrt{2} r} dt + \frac{g R Q_m}{2} \cos \theta_1 d\varphi_1.
\end{aligned} \tag{4.58}$$

For different values of the constants  $Q_e, Q_m, R, g$ , we will have either a horizon or a naked singularity. It should be instructive to compute the entropy and the Hawking temperature of this black hole and comparing them with those of the M-theory black holes calculated in [123]. Since the system analyzed in that reference is the same in string variables as the one we analyze here, it is expected that those results will be repeated. Indeed, the ten-dimensional interpretation suggested in [123] agrees with the one we have described above.

## 5. A solution with excited B-fields

Here we consider the six-dimensional Romans' theory with the mass parameter and all fields except the scalar and the 3-form field set to zero. In addition, we take 3-form  $G_3$  to be a constant. The geometry of the  $AdS_3 \times R^3$  space-time is given by

$$ds^2 = e^{2f} (dt^2 - dr^2 - dz^2) - e^{2h} (dx^2 + dy^2 + dv^2). \tag{4.59}$$

Let us consider a spinor satisfying the following constraints

$$\Gamma_2 \Gamma_7 \epsilon_i = \epsilon_i , \quad \Gamma_{456} \epsilon_i = \epsilon_i , \quad (4.60)$$

and

$$G_{xyv} = G = \text{constant} . \quad (4.61)$$

From the vanishing of the supersymmetric variation of gravitinos and gauginos we obtain two independent equations

$$h' = -e^f \left( \frac{g}{4\sqrt{2}} e^\varphi + \frac{G}{2} e^{-3h+2\varphi} \right) , \quad (4.62)$$

$$f' = -e^f \left( \frac{g}{4\sqrt{2}} e^\varphi - \frac{G}{2} e^{-3h+2\varphi} \right) , \quad (4.63)$$

$$\varphi' = -h' . \quad (4.64)$$

A fixed-point solution can be easily obtained

$$e^{\frac{\phi}{\sqrt{2}}} = \frac{g e^{3h}}{2\sqrt{2}G} . \quad (4.65)$$

The values of  $h(r)$  remain undetermined since the equations for  $\phi'$  and  $h'$  are proportional to each other. In this case,  $f(r)$  is given by

$$f(r) = -\log \left( \frac{g^2 e^{3h} r}{8G} \right) , \quad (4.66)$$

therefore the metric is given by

$$ds^2 = \left( \frac{8G}{g^2 e^{3h} r} \right)^2 (dt^2 - dr^2 - dz^2) - e^{2h} (dx^2 + dy^2 + dv^2) . \quad (4.67)$$

Eq. (4.64) implies that  $\varphi = -h$ , where, for simplicity, we omit integration constants.

The system of equations above can be solved leading to

$$f(h) = -h + \frac{1}{2} \log \Xi , \quad (4.68)$$

where

$$\Xi = \frac{g}{4\sqrt{2}} e^{4h} + \frac{G}{2} . \quad (4.69)$$

Next, if we make the change of the integration variable  $r \rightarrow h$ , such that

$$e^f dr = -\frac{e^{5h}}{\Xi} dh , \quad (4.70)$$

in terms of  $h$  the metric reads

$$ds^2 = -\frac{e^{10h}}{\Xi^2} dh^2 + e^{-2h} \Xi (dt^2 - dz^2) - e^{2h} (dx^2 + dy^2 + dv^2) . \quad (4.71)$$

Using Eq. (E.16) we can write down the field strength of the dual theory as follows

$$F_{trz} = -G e^{2\sqrt{2}\phi+3f-3h} . \quad (4.72)$$

The ten-dimensional solution using Eq. (2.67) is

$$\begin{aligned} ds_{10}^2 &= -\frac{1}{2} e^{\sqrt{2}\phi/4} ds_6^2 + e^{5\sqrt{2}\phi/4} dz_1^2 \\ &\quad + \frac{1}{g^2} e^{-3\sqrt{2}\phi/4} [d\theta^2 + \sin^2\theta d\varphi^2 + (d\psi + \cos\theta d\varphi)^2] , \\ F_4 &= -G e^{2\sqrt{2}\phi+3f-3h} dt \wedge dr \wedge dz \wedge dz_1 - \frac{1}{g^2} \sin\theta d\psi \wedge d\theta \wedge d\varphi \wedge dz_1 , \\ \hat{\phi} &= \frac{1}{\sqrt{2}} \phi . \end{aligned} \quad (4.73)$$

The metric has the form of a warped product of a six-dimensional space, a 3-sphere and a single coordinate. We also note that the  $G_3$  field does not play a rôle similar to the previous cases. In fact, there is no twisting in this case, since the space is not curved. One can view this solution as a NS5-brane of type IIB after wrapping one of the directions on a circle. It would be interesting to find an example in which the twisting on the curved manifold is performed by a  $B_2$  field.

### C. Supergravity duals of three-dimensional $\mathcal{N} = 1$ SYM theory on a torus

In this section, we concentrate on a system which, when up-lifted to ten dimensions, can be interpreted as type IIB NS5-branes wrapped on  $S^3 \times T^2$ . In particular, this  $S^3$  is embedded in a seven-dimensional manifold with  $G_2$  holonomy. We consider the decoupling limit of  $N$  NS5-branes wrapped on  $S^3 \times T^2$  [124], keeping the radii fixed. Since the brane worldvolume is curved, in order to define covariantly constant Killing spinors, the resulting field theory on the brane worldvolume will be twisted. In order to describe the flows between the six-dimensional field theory (defined in the NS5-brane worldvolume in the UV) and the three-dimensional field theory in the IR, we will start with the  $SO(4)$ -symmetric solution, obtained by [125] of the five-dimensional  $SU(2)$ -gauged  $\mathcal{N} = 4$  supergravity constructed by Romans [126]. That solution only contains magnetic non-Abelian and electric Abelian fields. The dual twisted field theory is defined on the NS5-brane worldvolume wrapped on  $S^3$ , whereas the other two spatial directions are wrapped on a torus. In the IR limit it corresponds to a three-dimensional twisted gauge field theory on  $\mathbb{R}^1 \times T^2$ , with 2 supercharges. It is worth noting that this theory does not come from an  $AdS_4$ -like manifold since its spatial directions do not live in the spatial sector of five-dimensional supergravity, but on the torus in the ten-dimensional theory. Several aspects of this theory as seen from the gauge field theory point of view, including some dual twisted gauge field theories, have been analyzed in [50]. We then up-lift the previously mentioned solution to massless type IIA supergravity on  $S^1 \times S^3$ .

If we turn off the electric Abelian fields, it is possible to find a solution for the five-dimensional gauged supergravity [125], which is singular. Using the criterion given in [39], one can see that the singularity of that solution is a “bad” type, so that in the IR, this solution does not represent a gauge field theory. Therefore, one may

say that the electric Abelian fields remove the singularity. It would be interesting to know whether the non-singular solution with non-vanishing Abelian 2-form is related to the rotation of the NS5-brane. If it were the case, it would probably be related to the mechanism studied in [127], leading to a de-singularization by rotation.

### 1. Supergravity duals of $\mathcal{N} = 1$ SYM theory in $D = 3$ on a torus

In this subsection, we study the supergravity dual of a three-dimensional  $\mathcal{N} = 1$  SYM theory on a torus [52]. The gravitational system we are dealing with can be understood as follows. Let us consider  $N$  type IIB NS5-branes. If the 5-branes were flat, the isometries of this system would be  $SO(1, 5) \times SO(4)$ . The first corresponds to the Lorentz group on the flat 5-brane worldvolumes, while the second one is the corresponding rotation group of the  $S^3$  transverse to the 5-brane directions. Since the NS5-branes are not flat but wrapped on a second  $S^3$  (in the five-dimensional Romans' theory), we have the following chain of breaking of the isometries  $SO(1, 9) \rightarrow SO(1, 5) \times SO(4) \rightarrow SO(4) \times SO(4)$ . There is also an additional isometry group corresponding to the torus, where the two additional spatial directions of the 5-brane are wrapped. On the other hand, the supergravity solution that we consider here has a global  $SO(4)$  symmetry, and its corresponding ansatz for the five-dimensional metric has the  $\mathbb{R}^1 \times S^3 \times \mathbb{R}^1$  geometry. The  $\mathbb{R}^1$ s correspond to the time and the radial coordinate, respectively. In ten dimensions, the solution has the geometry of the form  $(\mathbb{R}_0^1 \times S_{1,2,3}^3 \times \mathbb{R}_4^1) \times T_{5,6}^2 \times S_{7,8,9}^3$ , where the lower indices label the coordinates. Recall from the previous chapters that the seven-dimensional supergravity is related to the five-dimensional one through a  $T^2$  reduction, whereas the up-lifting to 10-dimensional theory is obtained through an  $S^3$ . In Table I, we schematically show the global structure of the ten-dimensional metric. The first five coordinates are arbitrarily chosen to represent the five-dimensional metric for the Romans' theory.

Table I. Structure of ten-dimensional metric

|                  |               |                  |             |               |
|------------------|---------------|------------------|-------------|---------------|
| 0                | 1 2 3         | 4                | 5 6         | 7 8 9         |
| $\mathbb{R}_0^1$ | $S_{1,2,3}^3$ | $\mathbb{R}_4^1$ | $T_{5,6}^2$ | $S_{7,8,9}^3$ |

From Table I, one can see that the NS5-brane is wrapped on the  $S^3$  (which belongs to the five-dimensional  $SU(2)$ -gauged supergravity metric ansatz), while its other two spatial directions are wrapped on  $T_{5,6}^2$ , i.e. the fifth and sixth directions.

Now, we focus on the twisting preserving 2 supercharges. As already mentioned above, there are three spatial directions of the NS5-branes wrapped on  $S^3$ . Therefore, the supersymmetry will be realized through a twisting. Also notice that the NS5-branes have two directions on a torus, so that these are not involved in a twisting. The brane worldvolume is on  $\mathbb{R}^1 \times S^3 \times T^2$ . The non-trivial part of the spin connection on this worldvolume is the  $SU(2)$  connection on the spin bundle of  $S^3$ . On the other hand, the normal bundle to the NS5-brane in the  $G_2$  manifold is given by  $SU(2) \times SU(2)$ , one of them being the spin bundle of  $S^3$ . In this case, the twisting consists in the identification of the  $SU(2)$  group of the spin bundle with one of the factors in the R-symmetry group of the 5-brane, i.e.  $SO(4)_R \rightarrow SU(2)_L \times SU(2)_R$ . It leads to a diagonal group  $SU(2)_D$ , so that it gives a twisted gauge theory. The resulting symmetry group is  $SO(1, 2) \times SU(2)_D \times SU(2)_R$ . In the UV limit, the global symmetry is  $SO(1, 5) \times SU(2)_L \times SU(2)_R$ , so that the four scalars transform as the representation  $(\mathbf{1}, \mathbf{2}, \mathbf{2})$  and there are also 16 supercharges. After the twisting we get 2 fermions (which are the 2 supercharges of the remaining unbroken supersymmetry) transforming in the  $(\mathbf{2}, \mathbf{1}, \mathbf{1})$  representation of  $SO(1, 2) \times SU(2)_D \times SU(2)_R$ . There are no scalars after twisting, while we get one vector field as it is before the twisting. Therefore,  $1/8$  of the supersymmetries are preserved, which is related to the fact that



the two  $S^3$ s, together with the radial coordinate are embedded in a  $G_2$  manifold. In this way, since our IR limit corresponds to setting the radial coordinate to be zero, it implies (as we will see) that the  $S^3$  part of the 5-brane will reduce to a point. This is in contrast with the fact that the transverse  $S^3$  and the torus get fixed radii. It shows that when one moves to the IR of the gauge theory (flowing in the radial coordinate on the gravity dual) three of the dimensions become very small and no low energy massless modes are excited on this two-space. Therefore, far in the IR the gauge theory is effectively three-dimensional.

In order to show explicitly how the theory flows to a three-dimensional SYM theory on a torus, we briefly describe the  $SO(4)$ -symmetric solution of the five-dimensional Romans' supergravity presented in [125]. Following [125], we consider a static field configuration, invariant under the  $SO(4)$  global symmetry group of spatial rotations. As already mentioned, the metric ansatz has the structure  $\mathbb{R} \times S^3 \times \mathbb{R}$  and it can be written as

$$ds_5^2 = e^{2\nu(r)} dt^2 - \frac{1}{M(r)} dr^2 - r^2 d\Omega_3^2, \quad (4.74)$$

where  $d\Omega_3^2$  is the metric on  $S^3$ . In terms of left-invariant 1-forms (3.4)-(3.5),  $d\Omega_3^2$  can be written as  $\sum_{i=1}^3 \sigma^i \sigma^i$ . We consider the non-Abelian gauge potential components written in terms of the left-invariant 1-forms

$$A^i = A_\mu^i dx^\mu = [w(r) + 1] \sigma^i, \quad (4.75)$$

so that they are invariant under the combined action of the  $SO(4)$  rotations and the  $SU(2)$  gauge transformations. The corresponding field strength is purely magnetic and is given by

$$F^i = dw \wedge \sigma^i - [w(r)^2 - 1] d\sigma^i, \quad (4.76)$$

while for the Abelian gauge potential we consider a purely electric ansatz

$$f(r) = Q(r) dt \wedge dr . \quad (4.77)$$

All other functions, i.e.  $\nu$ ,  $M$ ,  $w$ ,  $Q$  and the dilaton  $\phi$  are functions of the radial coordinate  $r$ . From the equation of motion of the dilaton, one gets

$$\nu(r) = \sqrt{\frac{2}{3}} (\phi(r) - \phi_0) , \quad (4.78)$$

where  $\phi_0$  is an integration constant. On the other hand, from the equation of motion of the Abelian field, we obtain

$$Q(r) = \frac{e^{5\nu(r)}}{r^3 \sqrt{M(r)}} [2w(r)^3 - 6w(r) + H] , \quad (4.79)$$

where  $H$  is an integration constant. Other equations of motion with the field configuration and the metric ansatz described above can be found in [125]. In order for the solution to preserve some supersymmetries, it must satisfy the equations obtained by setting to zero the supersymmetry transformations for gauginos and gravitinos Eqs. (F.3)-(F.4). After some algebra, we obtain the following first-order differential equations

$$\begin{aligned} M(r) &= \left( \frac{1}{3} \zeta^2 V - w \right)^2 + 2 \zeta^2 (w^2 - 1)^2 - \frac{2}{3} (w^2 - 1) + \frac{1}{18 \zeta^2} , \\ \frac{dw(r)}{d \log r} &= \frac{1}{6 \zeta^2 M} \left\{ -2 V (w^2 - 1) \zeta^4 + (H - 4 w^3) \zeta^2 - w \right\} , \\ \frac{d\zeta(r)}{d \log r} &= -\frac{\zeta}{3 M} [V^2 \zeta^4 + 12 \zeta^2 (w^2 - 1)^2 - 4 V w \zeta^2 + w^2 + 2] , \end{aligned} \quad (4.80)$$

where we have defined  $\zeta(r) = e^\nu / r$  and  $V(r) = 2w(r)^3 - 6w(r) + H$ . These equations are compatible with the equations of motion derived from the Romans' five-dimensional Lagrangian given in appendix F, and any solution of these first-order differential equations preserves two supersymmetries.

Since we are interested in the IR limit, i.e. when  $r \rightarrow 0$ , we obtain the expansions of the functions defining the metric, the magnetic non-Abelian and the electric Abelian fields for the five-dimensional ansatz. They are

$$\begin{aligned} w(r) &= 1 - \frac{1}{24} r^2 + \dots, \\ \zeta(r) &= \frac{1}{r} + \frac{7}{288} r + \dots, \\ M(r) &= 1 + \frac{5}{144} r^2 + \dots, \end{aligned} \tag{4.81}$$

and straightforwardly

$$\nu(r) = \frac{7}{288} r^2 + \dots, \tag{4.82}$$

while for the dilaton we obtain

$$\phi(r) = \phi_0 + \frac{7}{288} \sqrt{\frac{3}{2}} r^2 + \dots. \tag{4.83}$$

In this case we have taken  $H$  to be 4. Also, we get

$$Q(r) = \frac{1}{96} r + \frac{13}{13824} r^3 + \dots. \tag{4.84}$$

In this way, one can see that in the IR limit the non-Abelian gauge potential has a core, while both field strengths, i.e. the Abelian and the non-Abelian one are of the order  $r$  around  $r = 0$ . Furthermore, the above solution can be up-lifted, following the ansätze presented in chapter II and in chapter III, to either type IIA or type IIB theory. As we will see later, the IR limits in these two cases turn out to be the same.

## 2. Up-lifting to type IIB theory

First, we consider the case when the solution is up-lifted to type IIB supergravity.

From Eq.(3.12) the ten-dimensional metric is

$$\begin{aligned}
 d\hat{s}_{10}^2 &= -e^{\frac{13}{5\sqrt{6}}\phi} \left( e^{2\nu(r)} dt^2 - \frac{1}{M(r)} dr^2 - r^2 d\Omega_3^2 \right) \\
 &\quad + e^{\frac{3}{5\sqrt{6}}\phi} (dY^2 + dZ^2) + \frac{1}{4g^2} e^{-\frac{3}{\sqrt{6}}\phi} \sum_{i=1}^3 (\sigma^i - g A_1^i)^2, \\
 \hat{\phi} &= \sqrt{6} \phi.
 \end{aligned} \tag{4.85}$$

Using the previously calculated IR, expansion we can obtain the radii of the different manifolds. Thus, for the  $S^3$  involving the coordinates 1, 2 and 3, the radius is given by

$$R_{1,2,3}^2 = e^{13\phi_0/5\sqrt{6}} r^2 + \mathcal{O}(r^4). \tag{4.86}$$

On the other hand, the radii of  $T_{5,6}^2$ ,  $S_{7,8,9}^3$  are given by

$$\begin{aligned}
 R_T^2 &= e^{\frac{\sqrt{3}}{5\sqrt{2}}\phi_0} \left( 1 - \frac{7}{960} r^2 + \mathcal{O}(r^3) \right), \\
 R_{7,8,9}^2 &= \frac{1}{4g^2} e^{-\frac{3}{\sqrt{6}}\phi_0} \left( 1 - \frac{21}{576} r^2 + \mathcal{O}(r^3) \right).
 \end{aligned} \tag{4.87}$$

Without loss of generality, we can set  $\phi_0$  to zero. It is obvious from Eqs. (4.86) and (4.87) that in the limit  $r \rightarrow 0$ ,  $R_T$  and  $R_{7,8,9}$  remain finite, while  $R_{1,2,3} \rightarrow 0$ . Since the type IIB NS5-brane is wrapped on  $S_{1,2,3}^3$ ,  $T^2$ , and in the IR limit  $S_{1,2,3}^3$  effectively reduces to a point, in this limit we obtain a twisted gauge field theory defined on the torus.

### 3. Up-lifting to type IIA theory

Using the ansatz (2.68) after up-lifting to type IIA theory, the ten-dimensional solution is

$$\begin{aligned}
d\hat{s}_{10}^2 &= -e^{\frac{7}{8\sqrt{6}}\phi} \left( e^{2\nu(r)} dt^2 - \frac{1}{M(r)} dr^2 - r^2 d\Omega_3^2 \right) \\
&\quad + \frac{1}{4g^2} e^{-\frac{9}{8\sqrt{6}}\phi} \sum_{i=1}^3 \left( \sigma^i - g A_1^i \right)^2 + e^{-\frac{9}{8\sqrt{6}}\phi} dZ^2 + e^{\frac{15}{8\sqrt{6}}\phi} dY^2, \\
\hat{\phi} &= \frac{3}{4\sqrt{6}} \phi.
\end{aligned} \tag{4.88}$$

As we did for the type IIB case, we can obtain the radius

$$R_{1,2,3}^2 = e^{\frac{7}{8\sqrt{6}}\phi_0} r^2 + \mathcal{O}(r^4), \tag{4.89}$$

which shrinks to zero in the IR limit. In addition, the radii of  $S_5^1$ ,  $S_{6,7,8}^3$  and  $S_9^1$  are finite

$$\begin{aligned}
R_5^2 &= e^{\frac{15}{8\sqrt{6}}\phi_0} \left( 1 + \frac{105}{4608} r^2 + \mathcal{O}(r^3) \right), \\
R_{6,7,8}^2 &= \frac{1}{4g^2} e^{-\frac{9}{8\sqrt{6}}\phi_0} \left( 1 - \frac{27}{4608} r^2 + \mathcal{O}(r^3) \right), \\
R_9^2 &= e^{-\frac{9}{8\sqrt{6}}\phi_0} \left( 1 - \frac{27}{4608} r^2 + \mathcal{O}(r^3) \right).
\end{aligned} \tag{4.90}$$

Again, by considering  $\phi_0 = 0$ , in the IR, the radii  $R_5 = R_9$  and  $R_{6,7,8} = 1/(2g)$ , while  $R_{1,2,3} \rightarrow 0$ . Since the type IIA NS5-brane is wrapped on  $S_{1,2,3}^3$ ,  $S_5^1$  and  $S_9^1$ , and in the IR limit  $S_{1,2,3}^3$  effectively shrinks to a point as in the type IIB case, we get the same geometric reduction as in the previous case. Note that this can be obtained when  $\phi_0 = 0$ , so that the radii of the torus (in type IIB case) and the two  $S^1$ s (in type IIA case) are exactly the same.

In addition, in both cases, one can use the criterion for confinement given in [128, 129], in order to show that the corresponding static potential is confining.

#### 4. The singular $SO(4)$ -symmetric solution

A solution with no electric Abelian fields can be obtained by setting  $H$  to zero. It implies that  $w$ ,  $V$  and also  $Q$  are zero, as we expected since no electric field is excited. In this way, the first-order differential equations (4.80) can be easily integrated, yielding the relation

$$r = r_0 \frac{e^{1/24\zeta^2}}{\sqrt{\zeta}} , \quad (4.91)$$

where  $r_0$  is an integration constant. The metric is given by

$$ds_5^2 = r_0^2 e^{1/(12\zeta^2)} \left( \zeta dt^2 - \frac{1}{8\zeta^5} d\zeta^2 - \frac{1}{\zeta} d\Omega_3^2 \right) . \quad (4.92)$$

In addition, for the dilaton we have the following relation

$$e^{\sqrt{2}\phi/\sqrt{3}} = r_0 e^{1/24\zeta^2} \sqrt{\zeta} . \quad (4.93)$$

Using the criterion of [39], it is obvious that the IR singularity is not acceptable, in both type IIA and type IIB theories.

## CHAPTER V

## CONCLUSIONS

We have presented a number of reduction ansätze of type IIA, type IIB supergravities in ten dimensions, and eleven-dimensional supergravity on various spaces of the form  $S^m \times T^n$ . Cases that were presented are reductions of type IIA supergravity on  $S^3$ ,  $S^4$ ,  $S^1 \times S^3$  and  $S^1 \times S^3 \times S^1$  which give rise to seven-dimensional,  $SO(4)$ -gauged supergravity, six-dimensional,  $SO(5)$ -gauged supergravity, the dual  $SU(2)$ -gauged supergravity in  $D = 6$  and the Romans' theory without the  $U(1)$  gauge-coupling in  $D = 5$ , respectively; reductions of type IIB supergravity on  $S^5$  and  $S^3 \times T^2$ . Reduction of type IIB supergravity on one deformation of  $S^5$  gives rise to the  $\mathcal{N} = 4$ ,  $SU(2) \times U(1)$ -gauged supergravity in  $D = 5$  and on another deformation of  $S^5$  produces the subset of the fields of five-dimensional  $\mathcal{N} = 8$   $SO(6)$ -gauged supergravity, consisting of gravity, the fifteen  $SO(6)$  gauge fields, and the twenty scalars of the  $SL(6, \mathbb{R})/SO(6)$  submanifold of the full scalar manifold. This embedding can equivalently be viewed as a complete reduction ansatz (with no truncation of massless fields) for the ten-dimensional theory, comprising just gravity plus a self-dual 5-form, which itself is a consistent truncation of type IIB supergravity. In a full  $S^5$  reduction of type IIB supergravity there will be additional fields coming from the reduction of the NS-NS and R-R 2-form potentials, and from the dilaton and axion. A complete analysis of the  $S^5$  reduction can therefore be expected to be extremely complicated. In particular, for example, the **10** and  $\overline{\mathbf{10}}$  of pseudo-scalars lead to a considerably more complicated metric reduction ansatz. The ansatz for a subset of the fields that included one scalar and one pseudo-scalar was derived in [34], and in [130].

These cases presented here represent other elements in the accumulating body

of examples where “remarkable” Kaluza-Klein sphere reductions exist, even though there is no known group-theoretic explanation for their consistency. What is still lacking is a deeper understanding of why they should work. One might be tempted to think that supersymmetry could provide the key, but this evidently cannot in general be the answer, since there are some examples such as the consistent  $S^3$  and  $S^{D-3}$  reductions of the  $D$ -dimensional low-energy limit of the bosonic string in arbitrary dimension  $D$  that are obviously unrelated to supersymmetry [63].

Albeit we have no theoretical explanation of why the spherical reductions should work, knowing the non-linear ansätze has proven to be very useful in a number of contexts. We showed that using the non-linear Kaluza-Klein ansatz considerably simplifies the calculation of  $n$ -point correlation functions in AdS/CFT correspondence. Applying the method in [82] and using the non-linear ansatz, we derived the CFT operators which are in agreement with those obtained based on the conformal symmetry [18]. One implication of our result is to find the operators that deform CFT theory from supergravity solutions. This procedure is an exact opposite to what has been presented in the literature, where one first deforms the CFT by adding some operators, and then tries to find the corresponding supergravity solution via the AdS/CFT duality.

We also showed that non-linear Kaluza-Klein reduction ansätze are very useful in studying the supergravity duals of certain field theories. These investigations provide more examples to support the well-known AdS/CFT correspondence. In particular, we found certain supergravity solutions in the Romans’ gauged supergravities in  $D = 5$  and  $D = 6$ . Using the non-linear Kaluza-Klein ansätze, these solutions, after uplifting to type IIA, type IIB and massive type IIA supergravities, can be interpreted as a D4-brane and a NS5-brane wrapped on  $S^2$ ,  $S^3$  and  $S^3 \times T^2$ . These solutions are dual to certain twisted field theories. D4-brane wrapped on  $S^2$  is supergravity dual



to five-dimensional  $\mathcal{N} = 2$  SCFT, which in the IR flows to  $\mathcal{N} = 2$  SCFTs in  $D = 3$ . On the other hand, D4-brane wrapped on  $S^3$  is supergravity dual to  $\mathcal{N} = 2$  SCFT in  $D = 5$ , which flows to a two-dimensional  $(1, 1)$  CFT. We also found a solution that has “good” singularity in the sense of [39, 116]. After uplifting to type IIB, this solution is interpreted as NS5-brane wrapped on  $S^2$  which is dual to three-dimensional  $\mathcal{N} = 2$  super Yang-Mills theory. We also studied the NS5-brane of type IIB wrapped on  $S^3$ . This solution is dual to three-dimensional  $\mathcal{N} = 1$  SYM theory on a torus.

## REFERENCES

- [1] M.B. Green, J.H. Schwarz, E. Witten, Superstring Theory, vol. 1-2. Cambridge University Press, Cambridge, 1987.
- [2] J. Polchinski, String Theory, vol. 1-2. Cambridge University Press, Cambridge, 1998.
- [3] E. Cremmer, B. Julia, J. Scherk, Supergravity theory in 11 dimensions, Phys. Lett. B 76 (1978) 409–412.
- [4] J.H. Schwarz, Covariant field equations of chiral  $\mathcal{N} = 2$   $D = 10$  supergravity, Nucl. Phys. B 226 (1983) 269–288.
- [5] M.J. Duff, B.E.W. Nilsson, C.N. Pope, Kaluza-Klein supergravity, Phys. Rept. 130 (1986) 1–142.
- [6] A. Salam, E. Sezgin, Supergravities in Diverse Dimensions, vol. 1-2. North-Holland/World Scientific, Singapore, 1989.
- [7] E. Cremmer, B. Julia, H. Lü, C.N. Pope, Dualisation of dualities. I, Nucl. Phys. B 523 (1998) 73–144, hep-th/9710119.
- [8] E. Cremmer, I.V. Lavrinenko, H. Lü, C.N. Pope, K.S. Stelle, T.A. Tran, Euclidean-signature supergravities, dualities and instantons, Nucl. Phys. B 534 (1998) 40–82, hep-th/9803259.
- [9] C.M. Hull, B. Julia, Duality and moduli spaces for timelike reductions, Nucl. Phys. B 534 (1998) 250–260, hep-th/9803239.

- [10] M.J. Duff, C.N. Pope, Kaluza-Klein supergravity and the seven sphere, in Supersymmetry and Supergravity (S. Ferrara, J.G. Taylor, P. van Nieuwenhuizen, eds.), World Scientific, Singapore, 1983.
- [11] K. Pilch, P. van Nieuwenhuizen, P.K. Townsend, Compactification of  $D = 11$  supergravity on  $S^4$  (or  $11 = 7 + 4$ , too), Nucl. Phys. B 242 (1984) 377–392.
- [12] M. Günaydin, N. Marcus, The spectrum of the  $S^5$  compactification of the chiral  $\mathcal{N} = 2$ ,  $D = 10$  supergravity and the unitary supermultiplets of  $U(2, 2/4)$ , Class. Quantum Grav. 2 (1985) L11–L17.
- [13] H.J. Kim, L.J. Romans, P. van Nieuwenhuizen, The mass spectrum of chiral  $\mathcal{N} = 2$   $D = 10$  supergravity on  $S^5$ , Phys. Rev. D 32 (1985) 389–399.
- [14] C.N. Pope, K.S. Stelle, Zilch currents, supersymmetry and Kaluza-Klein consistency, Phys. Lett. B 198 (1987) 151–155.
- [15] B. de Wit, H. Nicolai, The consistency of the  $S^7$  truncation in  $D = 11$  supergravity, Nucl. Phys. B 281 (1987) 211–240.
- [16] J.M. Maldacena, The large  $N$  limit of superconformal field theories and supergravity, Adv. Theor. Math. Phys. 2 (1998) 231–252, hep-th/9711200.
- [17] S.S. Gubser, I.R. Klebanov, A.M. Polyakov, Gauge theory correlators from non-critical string theory, Phys. Lett. B 428 (1998) 105–114, hep-th/9802109.
- [18] E. Witten, Anti-de Sitter space and holography, Adv. Theor. Math. Phys. 2 (1998) 253–291, hep-th/9802150.
- [19] M. Cvetič, M.J. Duff, P. Hoxha, J.T. Liu, H. Lü, J.X. Lu, R. Martinez-Acosta, C.N. Pope, H. Sati, T.A. Tran, Embedding  $AdS$  black holes in ten and eleven dimensions, Nucl. Phys. B 558 (1999) 96–126, hep-th/9903214.

- [20] H. Nastase, D. Vaman, P. van Nieuwenhuizen, Consistent nonlinear KK reduction of 11D supergravity on  $AdS_7 \times S^4$  and self-duality in odd dimensions, Phys. Lett. B 469 (1999) 96–102, hep-th/9905075.
- [21] H. Nastase, D. Vaman, P. van Nieuwenhuizen, Consistency of the  $AdS_7 \times S^4$  reduction and the origin of self-duality in odd dimensions, Nucl. Phys. B 581 (2000) 179–239, hep-th/9911238.
- [22] M. Cvetič, H. Lü, C.N. Pope, Gauged six-dimensional supergravity from massive type IIA, Phys. Rev. Lett. 83 (1999) 5226–5229, hep-th/9906221.
- [23] M. Cvetič, H. Lü, C.N. Pope, J.F. Vazquez-Poritz, AdS in warped spacetimes, Phys. Rev. D 62 (2000) 122003, hep-th/0005246.
- [24] M. Cvetič, H. Lü, C.N. Pope, Consistent warped-space Kaluza-Klein reductions, half-maximal gauged supergravities and  $CP^n$  constructions, Nucl. Phys. B 597 (2001) 172–196, hep-th/0007109.
- [25] M. Cvetič, H. Lü, C.N. Pope, A. Sadrzadeh, T.A. Tran,  $S^3$  and  $S^4$  reductions of type IIA supergravity, Nucl. Phys. B 590 (2000) 233–251, hep-th/0005137.
- [26] L.J. Romans, The  $F(4)$  gauged supergravity in six dimensions, Nucl. Phys. B 269 (1986) 691–711.
- [27] H. Lü, C.N. Pope, Exact embedding of  $\mathcal{N} = 1, D = 7$  gauged supergravity in  $D = 11$ , Phys. Lett. B 467 (1999) 67–72, hep-th/9906168.
- [28] H. Lü, C.N. Pope, T.A. Tran, Five-dimensional  $\mathcal{N} = 4, SU(2) \times U(1)$  gauged supergravity from type IIB, Phys. Lett. B 475 (2000) 261–268, hep-th/9909203.
- [29] M. Cvetič, H. Lü, C.N. Pope, Four-dimensional  $\mathcal{N} = 4, SO(4)$  gauged supergravity from  $D = 11$ , Nucl. Phys. B 574 (2000) 761–781, hep-th/9910252.

- [30] H. Lu, C.N. Pope, Branes on the brane, Nucl. Phys. B 598 (2001) 492–508, hep-th/0008050.
- [31] M. Cvetič, H. Lü, C.N. Pope, Brane-world Kaluza-Klein reductions and branes on the brane, hep-th/0009183.
- [32] M. Cvetič, S.S. Gubser, H. Lü, C.N. Pope, Symmetric potentials of gauged supergravities in diverse dimensions and coulomb branch of gauge theories, Phys. Rev. D 62 (2000) 086003, hep-th/9909121.
- [33] M. Cvetič, H. Lü, C.N. Pope, A. Sadrzadeh, Consistency of Kaluza-Klein sphere reductions of symmetric potentials, Phys. Rev. D 62 (2000) 046005, hep-th/0002056.
- [34] M. Cvetič, H. Lü, C.N. Pope, Geometry of the embedding of supergravity scalar manifolds in  $D = 11$  and  $D = 10$ , Nucl. Phys. B 584 (2000) 149–170, hep-th/0002099.
- [35] M. Cvetič, H. Lü, C.N. Pope, A. Sadrzadeh, T.A. Tran, Consistent  $SO(6)$  reduction of type IIB supergravity on  $S^5$ , Nucl. Phys. B 586 (2000) 275–286, hep-th/0003103.
- [36] I.Y. Park, A. Sadrzadeh, T.A. Tran, Super Yang-Mills operators from the D3-brane action in a curved background, Phys. Lett. B 497 (2001) 303–308, hep-th/0010116.
- [37] S. Lee, S. Minwalla, M. Rangamani, N. Seiberg, Three-point functions of chiral operators in  $D = 4$ ,  $\mathcal{N} = 4$  SYM at large  $N$ , Adv. Theor. Math. Phys. 2 (1998) 697–718, hep-th/9806074.

- [38] M. Bershadsky, C. Vafa, V. Sadov, D-branes and topological field theories, Nucl. Phys. B 463 (1996) 420–434, hep-th/9511222.
- [39] J.M. Maldacena, C. Núñez, Supergravity description of field theories on curved manifolds and a no go theorem, Int. J. Mod. Phys. A 16 (2001) 822–855, hep-th/0007018.
- [40] J.M. Maldacena, C. Núñez, Towards the large  $N$  limit of pure  $\mathcal{N} = 1$  super Yang Mills, Phys. Rev. Lett. 86 (2001) 588–591, hep-th/0008001.
- [41] M. Alishahiha, Y. Oz, AdS/CFT and BPS strings in four dimensions, Phys. Lett. B 465 (1999) 136–141, hep-th/9907206.
- [42] A. Fayyazuddin, D.J. Smith, Localized intersections of M-fivebranes and four-dimensional superconformal field theories, JHEP 04 (1999) 030, hep-th/9902210.
- [43] A. Fayyazuddin, D.J. Smith, Warped AdS near-horizon geometry of completely localized intersections of M5-branes, JHEP 10 (2000) 023, hep-th/0006060.
- [44] B. Brinne, A. Fayyazuddin, S. Mukhopadhyay, D.J. Smith, Supergravity M5-branes wrapped on Riemann surfaces and their QFT duals, JHEP 12 (2000) 013, hep-th/0009047.
- [45] I.R. Klebanov, A.A. Tseytlin, Gravity duals of supersymmetric  $SU(N) \times SU(N+M)$  gauge theories, Nucl. Phys. B 578 (2000) 123–138, hep-th/0002159.
- [46] I.R. Klebanov, M.J. Strassler, Supergravity and a confining gauge theory: Duality cascades and  $\chi$ SB-resolution of naked singularities, JHEP 08 (2000) 052, hep-th/0007191.

- [47] B.S. Acharya, J.P. Gauntlett, N. Kim, Fivebranes wrapped on associative three-cycles, *Phys. Rev. D* 63 (2001) 106003, hep-th/0011190.
- [48] H. Nieder, Y. Oz, Supergravity and D-branes wrapping special lagrangian cycles, *JHEP* 03 (2001) 008, hep-th/0011288.
- [49] J.P. Gauntlett, N. Kim, D. Waldram, M-fivebranes wrapped on supersymmetric cycles, *Phys. Rev. D* 63 (2001) 126001, hep-th/0012195.
- [50] C. Núñez, I.Y. Park, M. Schvellinger, T.A. Tran, Supergravity duals of gauge theories from  $F(4)$  gauged supergravity in six dimensions, *JHEP* 04 (2001) 025, hep-th/0103080.
- [51] J.D. Edelstein, C. Núñez, D6-branes and M-theory geometrical transitions from gauged supergravity, *JHEP* 04 (2001) 028, hep-th/0103167.
- [52] M. Schvellinger, T.A. Tran, Supergravity duals of gauge field theories from  $SU(2) \times U(1)$  gauged supergravity in five dimensions, *JHEP* 06 (2001) 025, hep-th/0105019.
- [53] J.M. Maldacena, H. Nastase, The supergravity dual of a theory with dynamical supersymmetry breaking, hep-th/0105049.
- [54] J.P. Gauntlett, N. Kim, S. Pakis, D. Waldram, Membranes wrapped on holomorphic curves, hep-th/0105250.
- [55] R. Hernandez, Branes wrapped on coassociative cycles, hep-th/0106055.
- [56] J.P. Gauntlett, N. Kim, D. Martelli, D. Waldram, Wrapped fivebranes and  $\mathcal{N} = 2$  super Yang-Mills theory, hep-th/0106117.

- [57] J. Gomis, D-branes, holonomy and M-theory, Nucl. Phys. B 606 (2001) 3–17, hep-th/0103115.
- [58] J. Gomis, T. Mateos, D6-branes wrapping Kähler four-cycles, hep-th/0108080.
- [59] M. Pernici, K. Pilch, P. van Nieuwenhuizen, Gauged maximally extended supergravity in seven dimensions, Phys. Lett. B 143 (1984) 103–107.
- [60] A.H. Chamseddine, W.A. Sabra,  $D = 7$   $SU(2)$  gauged supergravity from  $D = 10$  supergravity, Phys. Lett. B 476 (2000) 415–419, hep-th/9911180.
- [61] H. Nastase, D. Vaman, On the nonlinear KK reductions on spheres of supergravity theories, Nucl. Phys. B 583 (2000) 211–236, hep-th/0002028.
- [62] M. Cvetič, J.T. Liu, H. Lü, C.N. Pope, Domain-wall supergravities from sphere reduction, Nucl. Phys. B 560 (1999) 230–256, hep-th/9905096.
- [63] M. Cvetič, H. Lü, C.N. Pope, Consistent Kaluza-Klein sphere reductions, Phys. Rev. D 62 (2000) 064028, hep-th/0003286.
- [64] A. Salam, J. Strathdee, On Kaluza-Klein theory, Annals Phys. 141 (1982) 316–352.
- [65] B. de Wit, H. Nicolai, N.P. Warner, The embedding of gauged  $\mathcal{N} = 8$  supergravity into  $D = 11$  supergravity, Nucl. Phys. B 255 (1985) 29–62.
- [66] B.E.W. Nilsson, On the embedding of  $D = 4$   $\mathcal{N} = 8$  gauged supergravity in  $D = 11$   $\mathcal{N} = 1$  supergravity, Phys. Lett. B 155 (1985) 54–58.
- [67] A. Khavaev, K. Pilch, N.P. Warner, New vacua of gauged  $\mathcal{N} = 8$  supergravity in five dimensions, Phys. Lett. B 487 (2000) 14–21, hep-th/9812035.



- [68] P.K. Townsend, P. van Nieuwenhuizen, Gauged seven-dimensional supergravity, Phys. Lett. B 125 (1983) 41–46.
- [69] A. Salam, E. Sezgin,  $SO(4)$  gauging of  $\mathcal{N} = 2$  supergravity in seven-dimensions, Phys. Lett. B 126 (1983) 295–300.
- [70] F. Giani, M. Pernici, P. van Nieuwenhuizen, Gauged  $\mathcal{N} = 4$   $D = 6$  supergravity, Phys. Rev. D 30 (1984) 1680–1687.
- [71] P.M. Cowdall, On gauged maximal supergravity in six dimensions, JHEP 06 (1999) 018, hep-th/9810041.
- [72] P.M. Cowdall, Supersymmetric electrovac in gauged supergravities, Class. Quantum Grav. 15 (1998) 2937–2953, hep-th/9710214.
- [73] M. Roček, E. Verlinde, Duality, quotients, and currents, Nucl. Phys. B 373 (1992) 630–646, hep-th/9110053.
- [74] X.C. de la Ossa, F. Quevedo, Duality symmetries from non-abelian isometries in string theory, Nucl. Phys. B 403 (1993) 377–394, hep-th/9210021.
- [75] E. Alvarez, L. Alvarez-Gaume, J.L.F. Barbon, Y. Lozano, Some global aspects of duality in string theory, Nucl. Phys. B 415 (1994) 71–100, hep-th/9309039.
- [76] E. Alvarez, L. Alvarez-Gaume, Y. Lozano, On non-abelian duality, Nucl. Phys. B 424 (1994) 155–183, hep-th/9403155.
- [77] E. Alvarez, L. Alvarez-Gaume, Y. Lozano, A canonical approach to duality transformations, Phys. Lett. B 336 (1994) 183–189, hep-th/9406206.
- [78] E. Alvarez, L. Alvarez-Gaume, Y. Lozano, An introduction to T duality in string theory, Nucl. Phys. Proc. Suppl. 41 (1995) 1–20, hep-th/9410237.

- [79] A. Bossard, M. Mohammadi, Non-abelian duality in the string effective action, Nucl. Phys. B 595 (2001) 93–118, hep-th/0008062.
- [80] M. Günaydin, L.J. Romans, N.P. Warner, Compact and noncompact gauged supergravity theories in five dimensions, Nucl. Phys. B 272 (1986) 598–646.
- [81] S. Ferrara, M.A. Lledo, A. Zaffaroni, Born-Infeld corrections to D3 brane action in  $AdS_5 \times S^5$  and  $\mathcal{N} = 4$ ,  $D = 4$  primary superfields, Phys. Rev. D 58 (1998) 105029, hep-th/9805082.
- [82] S.R. Das, S.P. Trivedi, Three-brane action and the correspondence between  $\mathcal{N} = 4$  Yang Mills theory and anti de Sitter space, Phys. Lett. B 445 (1998) 142–149, hep-th/9804149.
- [83] S.R. Das, S.D. Mathur, Comparing decay rates for black holes and D-branes, Nucl. Phys. B 478 (1996) 561–576, hep-th/9606185.
- [84] Jr. C.G. Callan, S.S. Gubser, I.R. Klebanov, A.A. Tseytlin, Absorption of fixed scalars and the D-brane approach to black holes, Nucl. Phys. B 489 (1997) 65–94, hep-th/9610172.
- [85] S. Ferrara, C. Fronsdal, A. Zaffaroni, On  $\mathcal{N} = 8$  supergravity on  $AdS_5$  and  $\mathcal{N} = 4$  superconformal Yang-Mills theory, Nucl. Phys. B 532 (1998) 153–162, hep-th/9802203.
- [86] H. Liu, A.A. Tseytlin,  $D = 4$  super Yang-Mills,  $D = 5$  gauged supergravity, and  $D = 4$  conformal supergravity, Nucl. Phys. B 533 (1998) 88–108, hep-th/9804083.
- [87] I.Y. Park, Fundamental versus solitonic description of D3-branes, Phys. Lett. B 468 (1999) 213–218, hep-th/9907142.

- [88] D.Z. Freedman, S.D. Mathur, A. Matusis, L. Rastelli, Correlation functions in the  $CFT_d/AdS_{d+1}$  correspondence, Nucl. Phys. B 546 (1999) 96–118, hep-th/9804058.
- [89] W. Mück, K.S. Viswanathan, Conformal field theory correlators from classical scalar field theory on  $AdS_{d+1}$ , Phys. Rev. D 58 (1998) 041901, hep-th/9804035.
- [90] M. Cederwall, A. von Gussich, B.E.W. Nilsson, A. Westerberg, The Dirichlet super-three-brane in ten-dimensional type IIB supergravity, Nucl. Phys. B 490 (1997) 163–178, hep-th/9610148.
- [91] E. Bergshoeff, P.K. Townsend, Super D-branes, Nucl. Phys. B 490 (1997) 145–162, hep-th/9611173.
- [92] M. Aganagic, J. Park, C. Popescu, J.H. Schwarz, Dual D-brane actions, Nucl. Phys. B 496 (1997) 215–230, hep-th/9702133.
- [93] R.R. Metsaev, A.A. Tseytlin, Supersymmetric D3-brane action in  $AdS_5 \times S^5$ , Phys. Lett. B 436 (1998) 281–288, hep-th/9806095.
- [94] R.R. Metsaev, A.A. Tseytlin, Type IIB superstring action in  $AdS_5 \times S^5$  background, Nucl. Phys. B 533 (1998) 109–126, hep-th/9805028.
- [95] R. Kallosh, J. Rahmfeld, The GS string action on  $AdS_5 \times S^5$ , Phys. Lett. B 443 (1998) 143–146, hep-th/9808038.
- [96] J. Hughes, J. Liu, J. Polchinski, Supermembranes, Phys. Lett. B 180 (1986) 370–374.
- [97] K. Becker, M. Becker, A. Strominger, Fivebranes, membranes and nonperturbative string theory, Nucl. Phys. B 456 (1995) 130–152, hep-th/9507158.

- [98] K. Intriligator, D.R. Morrison, N. Seiberg, Five-dimensional supersymmetric gauge theories and degenerations of Calabi-Yau spaces, Nucl. Phys. B 497 (1997) 56–100, hep-th/9702198.
- [99] S. Ferrara, A. Kehagias, H. Partouche, A. Zaffaroni,  $AdS_6$  interpretation of 5D superconformal field theories, Phys. Lett. B 431 (1998) 57–62, hep-th/9804006.
- [100] A. Brandhuber, Y. Oz, The D4-D8 brane system and five-dimensional fixed points, Phys. Lett. B 460 (1999) 307–312, hep-th/9905148.
- [101] O. Aharony, A. Hanany, K. Intriligator, N. Seiberg, M.J. Strassler, Aspects of  $\mathcal{N} = 2$  supersymmetric gauge theories in three dimensions, Nucl. Phys. B 499 (1997) 67–99, hep-th/9703110.
- [102] J. de Boer, K. Hori, H. Ooguri, Y. Oz, Mirror symmetry in three-dimensional gauge theories, quivers and D-branes, Nucl. Phys. B 493 (1997) 101–147, hep-th/9611063.
- [103] J. de Boer, K. Hori, H. Ooguri, Y. Oz, Z. Yin, Mirror symmetry in three-dimensional gauge theories,  $SL(2, Z)$  and D-brane moduli spaces, Nucl. Phys. B 493 (1997) 148–176, hep-th/9612131.
- [104] J. de Boer, K. Hori, Y. Oz, Dynamics of  $\mathcal{N} = 2$  supersymmetric gauge theories in three dimensions, Nucl. Phys. B 500 (1997) 163–191, hep-th/9703100.
- [105] J. de Boer, K. Hori, Y. Oz, Z. Yin, Branes and mirror symmetry in  $\mathcal{N} = 2$  supersymmetric gauge theories in three dimensions, Nucl. Phys. B 502 (1997) 107–124, hep-th/9702154.
- [106] E. Witten, Small instantons in string theory, Nucl. Phys. B 460 (1996) 541–559, hep-th/9511030.

- [107] J. Polchinski, E. Witten, Evidence for heterotic - type I string duality, Nucl. Phys. B 460 (1996) 525–540, hep-th/9510169.
- [108] J. Polchinski, S. Chaudhuri, C.V. Johnson, Notes on D-branes, hep-th/9602052.
- [109] J. Polchinski, TASI lectures on D-branes, hep-th/9611050.
- [110] N. Seiberg, Five-dimensional SUSY field theories, non-trivial fixed points and string dynamics, Phys. Lett. B 388 (1996) 753–760, hep-th/9608111.
- [111] D.R. Morrison, N. Seiberg, Extremal transitions and five-dimensional supersymmetric field theories, Nucl. Phys. B 483 (1997) 229–247, hep-th/9609070.
- [112] B. Janssen, P. Meessen, T. Ortin, The D8-brane tied up: String and brane solutions in massive type IIA supergravity, Phys. Lett. B 453 (1999) 229–236, hep-th/9901078.
- [113] M. Massar, J. Troost, D0-D8-F1 in massive IIA SUGRA, Phys. Lett. B 458 (1999) 283–287, hep-th/9901136.
- [114] F.A. Cachazo, C. Vafa, Type I' and real algebraic geometry, hep-th/0001029.
- [115] V. Balasubramanian, P. Kraus, A.E. Lawrence, Bulk vs. boundary dynamics in anti-de Sitter spacetime, Phys. Rev. D 59 (1999) 046003, hep-th/9805171.
- [116] S.S. Gubser, Curvature singularities: The good, the bad, and the naked, hep-th/0002160.
- [117] I.V. Lavrinenko, H. Lü, C.N. Pope, K.S. Stelle, Superdualities, brane tensions and massive IIA/IIB duality, Nucl. Phys. B 555 (1999) 201–227, hep-th/9903057.

- [118] A. Buchel, C.P. Herzog, I.R. Klebanov, L. Pando Zayas, A.A. Tseytlin, Non-extremal gravity duals for fractional D3-branes on the conifold, JHEP 04 (2001) 033, hep-th/0102105.
- [119] S.S. Gubser, C.P. Herzog, I.R. Klebanov, A.A. Tseytlin, Restoration of chiral symmetry: A supergravity perspective, JHEP 05 (2001) 028, hep-th/0102172.
- [120] A. Buchel, A. Frey, Comments on supergravity dual of pure  $\mathcal{N} = 1$  super Yang Mills theory with unbroken chiral symmetry, hep-th/0103022.
- [121] A.H. Chamseddine, M.S. Volkov, Non-Abelian BPS monopoles in  $\mathcal{N} = 4$  gauged supergravity, Phys. Rev. Lett. 79 (1997) 3343–3346, hep-th/9707176.
- [122] A.H. Chamseddine, M.S. Volkov, Non-Abelian solitons in  $\mathcal{N} = 4$  gauged supergravity and leading order string theory, Phys. Rev. D 57 (1998) 6242–6254, hep-th/9711181.
- [123] D. Klemm, BPS black holes in gauged  $\mathcal{N} = 4$ ,  $D = 4$  supergravity, Nucl. Phys. B 545 (1999) 461–478, hep-th/9810090.
- [124] N. Itzhaki, J.M. Maldacena, J. Sonnenschein, S. Yankielowicz, Supergravity and the large  $N$  limit of theories with sixteen supercharges, Phys. Rev. D 58 (1998) 046004, hep-th/9802042.
- [125] A.H. Chamseddine, M.S. Volkov, Non-abelian vacua in  $D = 5$ ,  $\mathcal{N} = 4$  gauged supergravity, JHEP 04 (2001) 023, hep-th/0101202.
- [126] L.J. Romans, Gauged  $\mathcal{N} = 4$  supergravities in five dimensions and their magnetovac backgrounds, Nucl. Phys. B 267 (1986) 433–447.
- [127] J.M. Maldacena, L. Maoz, De-singularization by rotation, hep-th/0012025.

- [128] Y. Kinar, E. Schreiber, J. Sonnenschein,  $Q\overline{Q}$  potential from strings in curved spacetime: Classical results, Nucl. Phys. B 566 (2000) 103–125, hep-th/9811192.
- [129] J. Sonnenschein, What does the string/gauge correspondence teach us about Wilson loops?, hep-th/0003032.
- [130] K. Pilch, N.P. Warner, A new supersymmetric compactification of chiral IIB supergravity, Phys. Lett. B 487 (2000) 22–29, hep-th/0002192.
- [131] W. Nahm, Supersymmetries and their representations, Nucl. Phys. B 135 (1978) 149–166.
- [132] M. Huq, M.A. Namazie, Kaluza-Klein supergravity in ten dimensions, Class. Quantum Grav. 2 (1985) 293–308.
- [133] E. Bergshoeff, C. Hull, T. Ortin, Duality in the type II superstring effective action, Nucl. Phys. B 451 (1995) 547–578, hep-th/9504081.
- [134] B.S. DeWitt, P. van Nieuwenhuizen, Explicit construction of the exceptional superalgebras  $F(4)$  and  $G(3)$ , J. Math. Phys. 23 (1982) 1953–1963.
- [135] R. D’Auria, S. Ferrara, S. Vaulà, Matter coupled  $F(4)$  supergravity and the  $AdS_6/CFT_5$  correspondence, JHEP 10 (2000) 013, hep-th/0006107.
- [136] L. Andrianopoli, R. D’Auria, S. Vaulà, Matter coupled  $F(4)$  gauged supergravity lagrangian, JHEP 05 (2001) 065, hep-th/0104155.
- [137] P.K. Townsend, K. Pilch, P. van Nieuwenhuizen, Selfduality in odd dimensions, Phys. Lett. B 136 (1984) 38–42.

## APPENDIX A

## ELEVEN-DIMENSIONAL SUPERGRAVITY

Eleven is a maximum space-time dimension in which a consistent supersymmetric theory can be constructed [131]. The  $\mathcal{N} = 1$  supergravity in eleven dimensions was constructed by Cremmer, Julia and Scherk [3]. The field content of eleven-dimensional supergravity consists of a graviton  $g_{MN}$ , a 3-form potential  $A_{(3)}$  whose field strength is  $F_{(4)} = dA_{(3)}$  (or in terms of indexes  $F_{MNPQ} = 4\partial_{[M}A_{NPQ]}$ ), and a gravitino  $\Psi_M$  with 44, 84 and 128 physical degrees of freedom. For a counting number degrees of freedom in various dimensions, see Table. II. The Lagrangian of the bosonic sector is

$$\begin{aligned} \mathcal{L} = & \sqrt{-g} \left( R - \frac{1}{2 \cdot 4!} F_{MNPQ} F^{MNPQ} \right) \\ & - \frac{1}{3!^2 \cdot 4!^2} \epsilon^{M_1 \dots M_{11}} F_{M_1 \dots M_4} F_{M_5 \dots M_8} A_{M_9 \dots M_{11}} , \end{aligned} \quad (\text{A.1})$$

where  $g = \det(g_{MN})$ . Eq. (A.1) can also be written in form notation

$$\mathcal{L}_{11} = R * \mathbb{1} - \frac{1}{2} * F_{(4)} \wedge F_{(4)} - \frac{1}{6} F_{(4)} \wedge F_{(4)} \wedge A_{(3)} . \quad (\text{A.2})$$

The supersymmetry transformation of gravitino in the absence of fermion is

$$\delta \Psi_M = \left[ \partial_M \Psi_N + \frac{1}{4} (\omega_{AB})_N \Gamma^{AB} \Psi_N \right] \epsilon - \frac{1}{288} (\Gamma_M^{PQRS} - 8 \delta_M^P \Gamma^{QRS}) F_{PQRS} \epsilon . \quad (\text{A.3})$$

The equations of motion of the eleven-dimensional supergravity are

$$R_{MN} = \frac{1}{12} \left( F_{MPQR} F_N^{PQR} - \frac{1}{12} g_{MN} F_{PQRS} F^{PQRS} \right) , \quad (\text{A.4})$$

and

$$d * F_{(4)} = \frac{1}{2} F_{(4)} \wedge F_{(4)} . \quad (\text{A.5})$$



Table II. On-shell degrees of freedom in  $D$  dimensions.  $\alpha = \frac{1}{2}D$  if  $D$  is even,  $\alpha = \frac{1}{2}(D - 1)$  if  $D$  is odd.

| Field                  | Notation                  | # degrees of freedom         |
|------------------------|---------------------------|------------------------------|
| $D$ -bein              | $e_M^A$                   | $\frac{D(D-3)}{2}$           |
| Majorana fermions      | $(\Psi_M)_{Maj}$          | $2^{(\alpha-1)}(D-3)$        |
| Majorana-Weyl fermions | $(\Psi_M)_{Maj-Weyl}$     | $2^{(\alpha-2)}(D-3)$        |
| $p$ -form              | $A_{M_1 M_2 \dots M_p}$   | $\frac{(D-2)!}{p!(D-2-p)!}$  |
| self-dual $p$ -form    | $A_{M_1 M_2 \dots M_p}^+$ | $\frac{(D-2)!}{2p!(D-2-p)!}$ |
| Majorana spinor        | $\chi_{Maj}$              | $2^{(\alpha-1)}$             |
| Majorana-Weyl spinor   | $\chi_{Maj-Weyl}$         | $2^{(\alpha-2)}$             |

## APPENDIX B

## TEN-DIMENSIONAL TYPE IIA SUPERGRAVITY

Ten-dimensional type IIA supergravity can be obtained by dimensional reducing eleven-dimensional supergravity on a circle [132]. The field content of type IIA supergravity comprises a vielbein  $e_M^A$ , a scalar  $\phi$ , an R-R 1-form potential  $\mathcal{A}_{(1)}$  whose field strength is  $\mathcal{F}_{(2)} = d\mathcal{A}_{(1)}$ , an NS-NS 2-form potential  $B_{(2)}$  whose field strength is  $F_{(3)} = dB_{(2)}$  and an R-R 3-form potential  $A_{(3)}$  whose field strength is  $F_{(4)} = dA_{(3)} + \mathcal{A}_{(1)} \wedge F_{(3)}$ , two gravitini, which have opposite chirality,  $\Psi_M^1$  and  $\Psi_M^2$ , and two dilatini  $\lambda_1$  and  $\lambda_2$ . The bosonic Lagrangian of the theory is

$$\begin{aligned} \mathcal{L}_{\text{IIA}} = & R * \mathbb{1} - \frac{1}{2} * d\phi \wedge d\phi - \frac{1}{2} e^{\frac{3}{2}\phi} * \mathcal{F}_{(2)} \wedge \mathcal{F}_{(2)} - \frac{1}{2} e^{\frac{1}{2}\phi} * F_{(4)} \wedge F_{(4)} \\ & - \frac{1}{2} e^{-\phi} * F_{(3)} \wedge F_{(3)} + \frac{1}{2} dA_{(3)} \wedge dA_{(3)} \wedge A_{(2)} , \end{aligned} \quad (\text{B.1})$$

where

$$F_{(4)} = dA_{(3)} - dA_{(2)} \wedge \mathcal{A}_{(1)} , \quad F_{(3)} = dA_{(2)} , \quad \mathcal{F}_{(2)} = d\mathcal{A}_{(1)} . \quad (\text{B.2})$$

In the absence of the fermionic background, the gravitino transformation rule is

$$\begin{aligned} \delta \Psi_M = & \left( \partial_M + \frac{1}{4} (\omega_{AB})_M \Gamma^{AB} \right) \epsilon + \frac{1}{64} e^{-\frac{3}{2}\phi} (\Gamma_M^{NP} - 14 \delta_M^N \Gamma^P) \Gamma^{11} F_{NP} \epsilon \\ & + \frac{1}{96} e^{\frac{1}{2}\phi} (\Gamma_M^{NPQ} - 9 \delta_M^N \Gamma^{PQ}) \Gamma^{11} F_{NPQ} \epsilon \\ & + \frac{i}{256} e^{-\frac{1}{4}\phi} (\Gamma_M^{NPQR} - \frac{20}{3} \delta_M^N \Gamma^{PQR}) F_{NPQR} \epsilon , \end{aligned} \quad (\text{B.3})$$

and the supersymmetry transformation of the dilatino is

$$\begin{aligned} \delta \lambda = & -\frac{\sqrt{2}}{4} \left( \partial_M + \frac{1}{4} (\omega_{AB})_M \Gamma^{AB} \right) \phi \Gamma^M \Gamma^{11} \epsilon + \frac{3}{16\sqrt{2}} e^{-\frac{3}{4}\phi} \Gamma^{MN} F_{MN} \epsilon \\ & + \frac{i}{24\sqrt{2}} e^{\frac{1}{2}\phi} \Gamma^{MNP} F_{MNP} \epsilon - \frac{i}{192\sqrt{2}} e^{\frac{1}{4}\phi} \Gamma^{MNPQ} F_{MNPQ} \epsilon , \end{aligned} \quad (\text{B.4})$$

where  $\Gamma^M$  are the ten-dimensional Dirac matrices and  $\Gamma^{11} = \text{i} \Gamma^0 \Gamma^1 \dots \Gamma^9$ .

The equations of motion of the Lagrangian (B.1) are

$$\begin{aligned}
d*d\phi &= \frac{1}{2} e^{-\phi} *F_{(3)} \wedge F_{(3)} - \frac{3}{4} e^{\frac{3}{2}\phi} *\mathcal{F}_{(2)} \wedge \mathcal{F}_{(2)} - \frac{1}{4} e^{\frac{1}{2}\phi} *F_{(4)} \wedge F_{(4)} , \\
d(e^{\frac{1}{2}\phi} *F_{(4)}) &= F_{(4)} \wedge F_{(3)} , \\
d(e^{\frac{3}{2}\phi} *\mathcal{F}_{(2)}) &= -e^{\frac{1}{2}\phi} *F_{(4)} \wedge F_{(3)} , \\
d(e^{-\phi} *F_{(3)}) &= \frac{1}{2} F_{(4)} \wedge F_{(4)} - e^{\frac{1}{2}\phi} *F_{(4)} \wedge \mathcal{F}_{(2)} .
\end{aligned} \tag{B.5}$$

## APPENDIX C

## TEN-DIMENSIONAL TYPE IIB SUPERGRAVITY

The field content of ten dimensional type IIB supergravity consists of the metric, a self-dual 5-form field strength, a scalar, an axion, an R-R 3-form, an NS-NS 3-form field strength, two gravitini, which have the same chirality,  $\Psi_M^1, \Psi_M^2$  and two dilatini  $\lambda_1, \lambda_2$ . There is no simple covariant Lagrangian for type IIB supergravity, on account of the self-duality constraint for the 5-form. However, one can write a Lagrangian in which the 5-form is unconstrained, which must then be accompanied by a self-duality condition which is imposed by hand at the level of the equations of motion [133]. The bosonic Lagrangian of this type IIB is

$$\begin{aligned} \mathcal{L}_{\text{IIB}} = & R * \mathbb{1} - \frac{1}{2} * d\phi \wedge d\phi - \frac{1}{2} e^{2\phi} * d\chi \wedge d\chi - \frac{1}{4} * H_{(5)} \wedge H_{(5)} \\ & - \frac{1}{2} e^{-\phi} * F_{(3)}^2 \wedge F_{(3)}^2 - \frac{1}{2} e^{\phi} * F_{(3)}^1 \wedge F_{(3)}^1 - \frac{1}{2} B_{(4)} \wedge dA_{(2)}^1 \wedge dA_{(2)}^2, \quad (\text{C.1}) \end{aligned}$$

where

$$\begin{aligned} F_{(3)}^2 &= dA_{(2)}^2, \quad F_{(3)}^1 = dA_{(2)}^1 - \chi dA_{(2)}^2, \\ H_{(5)} &= dB_{(4)} - \frac{1}{2} A_{(2)}^1 \wedge dA_{(2)}^2 + \frac{1}{2} A_{(2)}^2 \wedge dA_{(2)}^1. \quad (\text{C.2}) \end{aligned}$$

The equations of motion following from the Lagrangian, together with the self-duality condition, are

$$\begin{aligned} R_{MN} = & \frac{1}{2} \partial_M \phi \partial_N \phi + \frac{1}{2} e^{2\phi} \partial_M \chi \partial_N \chi + \frac{1}{96} H_{MN}^2 \\ & + \frac{1}{4} e^{\phi} \left( (F_{(3)}^1)_{MN}^2 - \frac{1}{12} (F_{(3)}^1)^2 g_{MN} \right) \\ & + \frac{1}{4} e^{-\phi} \left( (F_{(3)}^2)_{MN}^2 - \frac{1}{12} (F_{(3)}^2)^2 g_{MN} \right), \quad (\text{C.3}) \end{aligned}$$

$$d * d\phi = -e^{2\phi} * d\chi \wedge d\chi - \frac{1}{2} e^{\phi} * F_{(3)}^1 \wedge F_{(3)}^1 + \frac{1}{2} e^{-\phi} * F_{(3)}^2 \wedge F_{(3)}^2, \quad (\text{C.4})$$

$$d(e^{2\phi} * d\chi) = e^\phi * F_{(3)}^1 \wedge F_{(3)}^2, \quad (C.5)$$

$$d(e^\phi * F_{(3)}^1) = H_{(5)} \wedge F_{(3)}^2, \quad d(e^{-\phi} * F_{(3)}^2 - \chi e^\phi * F_{(3)}^1) = -H_{(5)} \wedge (F_{(3)}^1 + \chi F_{(3)}^2), \quad (C.6)$$

$$d(*H_{(5)}) = -F_{(3)}^1 \wedge F_{(3)}^2, \quad H_{(5)} = *H_{(5)}. \quad (C.7)$$

The equations (C.3)-(C.7) are precisely those which were found in [4].

In the absence of fermionic backgrounds and using complex notations the supersymmetry transformation for dilatino and gravitino are

$$\begin{aligned} \delta\Psi_M &= \mathcal{D}_M \epsilon + \frac{i}{480} \Gamma^{N_1 \dots N_5} \Gamma_M \epsilon F_{M_1 \dots M_5} + \frac{1}{96} (\Gamma_M^{NPQ} G_{NPQ} - 9 \Gamma^{NP} G_{MNP}) \epsilon^*, \\ \delta\lambda &= i \Gamma^M \epsilon^* P_M - \frac{i}{24} \Gamma^{MNP} \epsilon G_{MNP}, \end{aligned} \quad (C.8)$$

where

$$\begin{aligned} G_{MNP} &= f (F_{MNP} - B F_{MNP}^*), \quad F_{MNP} = 3 \partial_{[M} A_{NP]}, \\ F_{MNPQR} &= 5 \partial_{[M} A_{NPQR]} - \frac{5}{4} \text{Im} (A_{[MN} F_{PQR]}^*), \\ Q_M &= f^2 \text{Im} (B P_M^*), \quad P_M = f^2 \partial_M B, \quad f = \frac{1}{\sqrt{1 - |B|^2}}. \end{aligned} \quad (C.9)$$

In the above Eq. (C.9),  $B$  is a complex scalar. Using a simple field redefinition [133],  $B$  can be written in terms of dilaton  $\phi$  and axion  $\chi$  as

$$B = \frac{1 - e^{-2\phi} - \chi^2 + 2i\chi}{(1 + e^{-\phi})^2 + \chi^2}. \quad (C.10)$$

The relation between 4-form potentials of two notations is  $B_{MNPQ} = 4A_{MNPQ}$ . The complex 3-form field strength  $F_{(3)}$  can be written in terms of NS-NS 2-form potential  $A_{(2)}^2$  and R-R 2-form potential  $A_{(2)}^1$  as  $F_{MNP} = 3\partial_{[M} (A_{NP]}^2 - iA_{NP]}^1)$ .

## APPENDIX D

## TEN-DIMENSIONAL MASSIVE TYPE IIA SUPERGRAVITY

The bosonic sector of massive type IIA supergravity [26] consists of a graviton, a scalar, a 2-form field strength, 3-form and 4-form field strengths. In the language of differential forms, the bosonic Lagrangian is given by [117]

$$\begin{aligned}
\mathcal{L}_{\text{massive}} = & R * \mathbb{1} - \frac{1}{2} * d\phi \wedge d\phi - \frac{1}{2} e^{\frac{3}{2}\phi} * F_{(2)} \wedge F_{(2)} - \frac{1}{2} e^{-\phi} * F_{(3)} \wedge F_{(3)} \\
& - \frac{1}{2} e^{\frac{1}{2}\phi} * F_{(4)} \wedge F_{(4)} - \frac{1}{2} dA_{(3)} \wedge dA_{(3)} \wedge A_{(2)} - \frac{1}{6} m dA_{(3)} \wedge (A_{(2)})^3 \\
& - \frac{1}{40} m^2 (A_{(2)})^5 - \frac{1}{2} m^2 e^{\frac{5}{2}\phi} * \mathbb{1} ,
\end{aligned} \tag{D.1}$$

where the field strengths are given in terms of potentials by

$$\begin{aligned}
F_{(2)} &= dA_{(1)} + m A_{(2)} , & F_{(3)} &= dA_{(2)} , \\
F_{(4)} &= dA_{(3)} + A_{(1)} \wedge dA_{(2)} + \frac{1}{2} m A_{(2)} \wedge A_{(2)} .
\end{aligned} \tag{D.2}$$

It follows that the equations of motion and Bianchi identities are

$$\begin{aligned}
d(e^{\frac{1}{2}\phi} * F_{(4)}) &= -F_{(3)} \wedge F_{(4)} , & d(e^{\frac{3}{2}\phi} * F_{(2)}) &= -e^{\frac{1}{2}\phi} * F_{(4)} \wedge F_{(3)} , \\
d(e^{-\phi} * F_{(3)}) &= -\frac{1}{2} F_{(4)} \wedge F_{(4)} - m e^{\frac{3}{2}\phi} * F_{(2)} - e^{\frac{1}{2}\phi} * F_{(4)} \wedge F_{(2)} , \\
d * d\phi &= -\frac{5}{4} m^2 e^{\frac{5}{2}\phi} * \mathbb{1} - \frac{3}{4} e^{\frac{3}{2}\phi} * F_{(2)} \wedge F_{(2)} + \frac{1}{2} e^{-\phi} * F_{(3)} \wedge F_{(3)} - \frac{1}{4} e^{\frac{1}{2}\phi} * F_{(4)} \wedge F_{(4)} , \\
dF_{(4)} &= F_{(2)} \wedge F_{(3)} , & dF_{(3)} &= 0 , & dF_{(2)} &= m F_{(3)} ,
\end{aligned} \tag{D.3}$$

for the form fields and dilaton, together with the Einstein equation (in vielbein components)

$$\begin{aligned}
R_{AB} = & \frac{1}{2} \partial_A \phi \partial_B \phi + \frac{1}{16} m^2 e^{\frac{5}{2}\phi} \eta_{AB} + \frac{1}{12} e^{\frac{1}{2}\phi} \left( F_{(4)AB}^2 - \frac{3}{32} F_{(4)}^2 \eta_{AB} \right) \\
& + \frac{1}{4} e^{-\phi} \left( F_{(3)AB}^2 - \frac{1}{12} F_{(3)}^2 \eta_{AB} \right) + \frac{1}{2} e^{\frac{3}{2}\phi} \left( F_{(2)AB}^2 - \frac{1}{16} F_{(2)}^2 \eta_{AB} \right) .
\end{aligned} \tag{D.4}$$

## APPENDIX E

## SIX-DIMENSIONAL GAUGED SUPERGRAVITY

The existence of six-dimensional gauged supergravity possessing a ground state with unbroken  $F_4$  supersymmetry was predicted by Nahm [131] and by DeWitt and van Nieuwenhuizen [134]. The six-dimensional gauged  $\mathcal{N} = 4$  supergravity constructed by Romans [26]. Further developments of  $F_4$  gauged supergravity in six dimensions coupled to matter were recently done by D'Auria et al. [135, 136]. Here we will review the theory constructed by Romans [26], whose conventions we follow. The theory consists of a graviton  $e_\mu^\alpha$ , three  $SU(2)$  gauge potentials  $A_\mu^I$ , an Abelian potential  $\mathcal{A}_\mu$ , a two-index tensor gauge field  $B_{\mu\nu}$ , a scalar  $\phi$ , four gravitinos  $\psi_{\mu i}$  and four gauginos  $\chi_\mu$ . The bosonic Lagrangian is

$$\begin{aligned}
e^{-1} \mathcal{L} = & -\frac{1}{4} R + \frac{1}{2} (\partial^\mu \phi) (\partial_\mu \phi) - \frac{1}{4} e^{-\sqrt{2}\phi} (\mathcal{H}^{\mu\nu} \mathcal{H}_{\mu\nu} + F^{I\mu\nu} F_{\mu\nu}^I) \\
& + \frac{1}{12} e^{2\sqrt{2}\phi} G_{\mu\nu\rho} G^{\mu\nu\rho} + \frac{1}{8} (g^2 e^{\sqrt{2}\phi} + 4g m e^{-\sqrt{2}\phi} - m^2 e^{-3\sqrt{2}\phi}) \\
& - \frac{1}{8} e^{-1} \varepsilon^{\mu\nu\rho\sigma\tau\kappa} B_{\mu\nu} (\mathcal{F}_{\rho\sigma} \mathcal{F}_{\tau\kappa} + m B_{\rho\sigma} \mathcal{F}_{\tau\kappa} \\
& + \frac{1}{3} m^2 B_{\rho\sigma} B_{\tau\kappa} + F_{\rho\sigma}^I F_{\tau\kappa}^I) , \tag{E.1}
\end{aligned}$$

where  $e$  is the determinant of the vielbein,  $g$  is the  $SU(2)$  coupling constant,  $m$  is the mass parameter associated with the two-index tensor field  $B_{\mu\nu}$  and  $\varepsilon_{\mu\nu\rho\sigma\tau\kappa}$  is a Levi-Civita tensor density. The Abelian field strength  $\mathcal{F}_{\mu\nu}$ , the non-Abelian one  $F_{\mu\nu}^I$ , the three-form  $G_{\mu\nu\rho}$  and the field  $\mathcal{H}_{\mu\nu}$  are given by

$$\begin{aligned}
\mathcal{F}_{\mu\nu} & \equiv \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu , \quad F_{\mu\nu}^I \equiv \partial_\mu A_\nu^I - \partial_\nu A_\mu^I + g \epsilon^{IJK} A_\mu^J A_\nu^K , \\
G_{\mu\nu\rho} & \equiv 3 \partial_{[\mu} B_{\nu\rho]} , \quad \mathcal{H}_{\mu\nu} \equiv \mathcal{F}_{\mu\nu} + m B_{\mu\nu} , \tag{E.2}
\end{aligned}$$

respectively. The supersymmetry transformations for the gauginos is

$$\delta\chi_i = \left( \frac{1}{\sqrt{2}} \Gamma^\mu \partial_\mu \phi + A \Gamma_7 - \frac{1}{12} e^{\sqrt{2}\phi} \Gamma_7 \Gamma^{\mu\nu\rho} G_{\mu\nu\rho} \right) \epsilon_i + \frac{1}{2\sqrt{2}} \Gamma^{\mu\nu} (\hat{H}_{\mu\nu})_i{}^j \epsilon_j, \quad (\text{E.3})$$

and that for the gravitinos is

$$\begin{aligned} \delta\psi_{\mu i} = & \left( \nabla_\mu + T \Gamma_\mu \Gamma_7 - \frac{1}{24} e^{\sqrt{2}\phi} \Gamma_7 \Gamma^{\nu\rho\sigma} G_{\nu\rho\sigma} \Gamma_\mu \right) \epsilon_i \\ & + \left[ g A_\mu^I (T^I)_i{}^j - \frac{1}{4\sqrt{2}} (\Gamma_\mu{}^{\nu\rho} - 6 \delta_\mu{}^\nu \Gamma^\rho) (\hat{H}_{\nu\rho})_i{}^j \right] \epsilon_j, \end{aligned} \quad (\text{E.4})$$

where  $A$ ,  $T$ , and  $\hat{H}$  are defined as follows

$$A \equiv \frac{1}{4\sqrt{2}} (g e^{\frac{1}{\sqrt{2}}\phi} - 3 m e^{\frac{-3}{\sqrt{2}}\phi}), \quad T \equiv -\frac{1}{8\sqrt{2}} (g e^{\frac{1}{\sqrt{2}}\phi} + m e^{\frac{-3}{\sqrt{2}}\phi}), \quad (\text{E.5})$$

$$(\hat{H}_{\mu\nu})_i{}^j \equiv e^{-\frac{1}{\sqrt{2}}\phi} \left[ \frac{1}{2} \mathcal{H}_{\mu\nu} \delta_i{}^j + \Gamma_7 F_{\mu\nu}^I (T^I)_i{}^j \right]. \quad (\text{E.6})$$

The gauge-covariant derivative  $\mathcal{D}_\mu$  acting on the Killing spinor is

$$\mathcal{D}_\mu \epsilon_i = \nabla_\mu \epsilon_i + g A_\mu^I (T^I)_i{}^j \epsilon_j, \quad (\text{E.7})$$

with

$$\nabla_\mu \epsilon_i \equiv \left( \partial_\mu + \frac{1}{4} \omega_\mu{}^{\alpha\beta} \Gamma_{\alpha\beta} \right) \epsilon_i, \quad (\text{E.8})$$

where  $\omega_\mu{}^{\alpha\beta}$  is the spin connection. Indices  $\alpha$  and  $\beta$  are tangent space (or flat) indices, while  $\mu$  and  $\nu$  are space-time (or curved) indices. The  $\Gamma_{\alpha\beta\dots}$  are the six-dimensional Dirac matrices,

$$\Gamma_{\alpha_1 \dots \alpha_n} = \frac{1}{n!} \Gamma_{[\alpha_1} \dots \Gamma_{\alpha_n]}, \quad n = 1, \dots, 6.$$

The Einstein equation of the Lagrangian (E.1) is

$$\begin{aligned} R_{\mu\nu} = & 2 \partial_\mu \phi \partial_\nu \phi + \frac{1}{8} g_{\mu\nu} (g^2 e^{\sqrt{2}\phi} + 4 g m e^{-\sqrt{2}\phi} - m^2 e^{-3\sqrt{2}\phi}) \\ & + e^{2\sqrt{2}\phi} \left( G_\mu{}^{\rho\sigma} G_{\nu\rho\sigma} - \frac{1}{6} g_{\mu\nu} G^{\rho\sigma\tau} G_{\rho\sigma\tau} \right) \end{aligned}$$



$$\begin{aligned}
& -2 e^{-\sqrt{2}\phi} \left( \mathcal{H}_\mu{}^\rho \mathcal{H}_{\nu\rho} - \frac{1}{8} g_{\mu\nu} \mathcal{H}_{\rho\sigma} \mathcal{H}^{\rho\sigma} \right) \\
& -2 e^{-\sqrt{2}\phi} \left( F_\mu{}^\rho F_{\nu\rho}^I - \frac{1}{8} g_{\mu\nu} F_{\rho\sigma}^I F^{I\rho\sigma} \right) .
\end{aligned} \tag{E.9}$$

The dilaton equation is

$$\begin{aligned}
\Box\phi &= \frac{1}{4\sqrt{2}} (g^2 e^{\sqrt{2}\phi} - 4 m g e^{-\sqrt{2}\phi} + 3 m^2 e^{-3\sqrt{2}\phi}) + \frac{1}{3\sqrt{2}} e^{2\sqrt{2}\phi} G^{\mu\nu\rho} G_{\mu\nu\rho} \\
&+ \frac{1}{2\sqrt{2}} e^{-\sqrt{2}\phi} (\mathcal{H}^{\mu\nu} \mathcal{H}_{\mu\nu} + F^{I\mu\nu} F_{\mu\nu}^I) ,
\end{aligned} \tag{E.10}$$

and the equations of motion for gauge fields are

$$\mathcal{D}_\nu (e^{-\sqrt{2}\phi} \mathcal{H}^{\nu\mu}) = \frac{1}{6} e \varepsilon^{\mu\nu\rho\sigma\tau\kappa} \mathcal{H}_{\nu\rho} G_{\sigma\tau\kappa} , \tag{E.11}$$

$$\mathcal{D}_\nu (e^{-\sqrt{2}\phi} F^{I\nu\mu}) = \frac{1}{6} e \varepsilon^{\mu\nu\rho\sigma\tau\kappa} F_{\nu\rho}^I G_{\sigma\tau\kappa} , \tag{E.12}$$

$$\mathcal{D}_\rho (e^{2\sqrt{2}\phi} G^{\rho\mu\nu}) = -m e^{-\sqrt{2}\phi} \mathcal{H}^{\mu\nu} - \frac{1}{4} e \varepsilon^{\mu\nu\rho\sigma\tau\kappa} (\mathcal{H}_{\rho\sigma} \mathcal{H}_{\tau\kappa} + F_{\rho\sigma}^I F_{\tau\kappa}^I) . \tag{E.13}$$

Depending upon the values of the gauge coupling and mass parameter, there are five distinct theories:  $\mathcal{N} = 4^+$  (for  $g > 0, m > 0$ ),  $\mathcal{N} = 4^-$  (for  $g < 0, m > 0$ ),  $\mathcal{N} = 4^g$  (for  $g > 0, m = 0$ ),  $\mathcal{N} = 4^m$  (for  $g = 0, m > 0$ ), and  $\mathcal{N} = 4^0$  (for  $g = 0, m = 0$ ). The  $\mathcal{N} = 4^g$  theory coincides with a theory [70] obtained by dimensional reduction of gauged  $\mathcal{N} = 2$  supergravity in seven dimensions, followed by truncation. It was pointed out in [26] that there is a dual version of the  $\mathcal{N} = 4^g$  theory, which has a similar field content but cannot be obtained from the  $\mathcal{N} = 4^g$  theory by a field redefinition. The  $B_{\mu\nu}$  field is replaced by a new tensor field  $A_{\mu\nu}$ , whose field strength is  $\tilde{F}_{\mu\nu}^I$ . In absence of the Abelian field strength, the bosonic Lagrangian of the dual  $\mathcal{N} = \tilde{4}^g$  theory is

$$\begin{aligned}
\tilde{e}^{-1} \tilde{\mathcal{L}} &= -\frac{1}{4} \tilde{R} + \frac{1}{2} \partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi} - \frac{1}{4} e^{-\sqrt{2}\tilde{\phi}} \tilde{F}_{\mu\nu}^I \tilde{F}^{I\mu\nu} \\
&+ \frac{1}{12} e^{-2\sqrt{2}\tilde{\phi}} \tilde{F}_{\mu\nu\rho} \tilde{F}^{\mu\nu\rho} + \frac{1}{8} \tilde{g}^2 e^{\sqrt{2}\tilde{\phi}} ,
\end{aligned} \tag{E.14}$$

where

$$\tilde{F}_{\mu\nu\rho} \equiv 3 \left( \partial_{[\mu} A_{\nu\rho]} - F_{[\mu\nu}^I A_{\rho]}^I - \frac{1}{3} g \epsilon^{IJK} A_\mu^I A_\nu^J A_\rho^K \right). \quad (\text{E.15})$$

The Lagrangian (E.14) can be obtained from the Lagrangian (E.1) by formally writing  $\tilde{F}_{\mu\nu\rho}$  as follows

$$\tilde{F}_{\mu\nu\rho} = \frac{1}{6} e^{2\sqrt{2}\tilde{\phi}} e \varepsilon_{\mu\nu\rho\sigma\tau\kappa} G^{\sigma\tau\kappa}. \quad (\text{E.16})$$

Since the Lagrangian (E.14) differs from  $\mathcal{N} = 4^g$  theory only by a sign of dilaton coupling to three-form field, in the absence of the three-form field  $\mathcal{N} = 4^g$  and  $\mathcal{N} = \tilde{4}^g$  are identical. Furthermore, the Eq. (E.16) tells us that any solution of  $\mathcal{N} = 4^g$  theory with the two-form fields not being excited, which is also a solution of  $\mathcal{N} = \tilde{4}^g$  theory, can be up-lifted into higher dimensional theory.

After re-scaling, the Lagrangian (E.1) can be written in terms of form notation

$$\begin{aligned} \mathcal{L} = & R * \mathbb{1} - \frac{1}{2} * d\phi \wedge d\phi - \frac{1}{2} e^{\frac{1}{\sqrt{2}}\phi} (*\mathcal{H}_2 \wedge \mathcal{H}_{(2)} + *F_{(2)}^I \wedge F_{(2)}^i) \\ & - \frac{1}{2} e^{-\sqrt{2}\phi} *G_{(3)} \wedge G_{(3)} + \frac{1}{16} (4g^2 e^{-\frac{1}{\sqrt{2}}\phi} + 2mg e^{\frac{1}{\sqrt{2}}\phi} - m^2 e^{-\frac{3}{\sqrt{2}}\phi}) * \mathbb{1} \\ & - B_{(2)} \wedge \left( \frac{1}{2} \mathcal{F}_{(2)} \wedge \mathcal{F}_{(2)} \right. \\ & \left. + \frac{1}{2} m B_{(2)} \wedge \mathcal{F}_{(2)} + \frac{1}{3} m^2 B_{(2)} \wedge B_{(2)} + \frac{1}{2} F_{(2)}^I \wedge F_{(2)}^I \right). \end{aligned} \quad (\text{E.17})$$

The dilaton equation in terms of form notation is

$$\begin{aligned} d*d\phi = & \frac{1}{\sqrt{2}} e^{-\sqrt{2}\phi} *G_{(3)} \wedge G_{(3)} - \frac{1}{2\sqrt{2}} e^{\frac{1}{\sqrt{2}}\phi} (*\mathcal{H}_{(2)} \wedge \mathcal{H}_{(2)} + *F_{(2)}^I \wedge F_{(2)}^I) \\ & - \frac{1}{16} \left( \frac{3}{\sqrt{2}} m^2 e^{-\frac{3}{\sqrt{2}}\phi} + \sqrt{2} m g e^{\frac{1}{\sqrt{2}}\phi} - 2\sqrt{2} g^2 e^{-\frac{1}{\sqrt{2}}\phi} \right) * \mathbb{1}. \end{aligned} \quad (\text{E.18})$$

the equations of motion for gauge fields are

$$\begin{aligned} d(e^{-\sqrt{2}\phi} *G_{(3)}) &= -\frac{1}{2} \mathcal{H}_{(2)} \wedge \mathcal{H}_{(2)} - \frac{1}{2} F_{(2)}^I \wedge F_{(2)}^I - m e^{\frac{1}{\sqrt{2}}\phi} *\mathcal{H}_{(2)}, \\ d(e^{\frac{1}{\sqrt{2}}\phi} *\mathcal{H}_{(2)}) &= -G_{(3)} \wedge \mathcal{H}_{(2)}, \end{aligned}$$

$$\mathcal{D}(e^{-\frac{1}{\sqrt{2}}\phi} * F_{(2)}^I) = -F_{(2)}^I \wedge G_{(3)}, \quad (\text{E.19})$$

and the Einstein equation is

$$\begin{aligned} R_{\mu\nu} = & -\frac{1}{2} \partial_\mu \phi \partial_\nu \phi + \left( \frac{1}{8} m^2 e^{\frac{3}{\sqrt{2}}\phi} - m g e^{\frac{1}{\sqrt{2}}\phi} - \frac{1}{2} g^2 e^{-\frac{1}{\sqrt{2}}\phi} \right) g_{\mu\nu} \\ & + \frac{1}{4} e^{-\sqrt{2}\phi} \left( G_{\mu\rho\sigma} G_\nu{}^{\rho\sigma} - \frac{1}{6} g_{\mu\nu} G_{\rho\sigma\tau} G^{\rho\sigma\tau} \right) \\ & + \frac{1}{2} e^{\frac{1}{\sqrt{2}}\phi} \left( \mathcal{H}_{\mu\rho} \mathcal{H}_\nu{}^\rho - \frac{1}{8} g_{\mu\nu} \mathcal{H}_{\rho\sigma} \mathcal{H}^{\rho\sigma} \right) \\ & + \frac{1}{2} e^{\frac{1}{\sqrt{2}}\phi} \left( F_{\mu\rho}^I F_\nu{}^{\rho\sigma} - \frac{1}{8} g_{\mu\nu} F_{\rho\sigma}^I F^{\rho\sigma} \right). \end{aligned} \quad (\text{E.20})$$

As shown in [22], the six-dimensional Romans' theory (E.17) can be obtained by reducing massive type IIA on a local  $S^4$ . We present the results which are written in terms of two parameters  $m$  and  $g$ . The metric ansatz is

$$\begin{aligned} d\hat{s}_{10}^2 = & \left( \frac{1}{3mg^2} \right)^{1/8} (\sin \xi)^{1/12} X^{1/8} \left[ -\frac{1}{2} \Delta^{3/8} ds_6^2 + g^2 \Delta^{3/8} X^2 d\xi^2 \right. \\ & \left. + \frac{1}{g} \Delta^{-5/8} X^{-1} \cos^2 \xi \sum_{i=1}^3 (\sigma^i - g A_{(1)}^i)^2 \right]. \end{aligned} \quad (\text{E.21})$$

The ansätze for 4-form, 3-form and 2-form are

$$\begin{aligned} \hat{F}_{(4)} = & \left( \frac{1}{3mg^3} \right)^{3/4} \left[ -\frac{2}{3} s^{1/3} c^3 \Delta^{-2} U d\xi \wedge \epsilon_{(3)} - 12mg s^{4/3} c^4 \Delta^{-2} X^{-3} dX \wedge \epsilon_{(3)} \right. \\ & + 2g^2 c s^{1/3} X^4 * G_{(3)} \wedge d\xi - \frac{3mg^2}{2} s^{4/3} X^{-2} * \mathcal{H}_{(2)} \\ & \left. + 2g c s^{1/3} F_{(2)}^i \wedge h^i \wedge d\xi - \frac{3mg}{2} c^2 s^{4/3} \Delta^{-1} X^{-3} F_{(2)}^i \wedge h^j \wedge h^k \epsilon_{ijk} \right], \\ \hat{F}_{(2)} = & \left( \frac{2g}{3m} \right)^{1/4} s^{2/3} \mathcal{H}_{(2)}, \\ \hat{F}_{(3)} = & \left( \frac{1}{3mg} \right)^{1/2} [3m s^{2/3} F_{(3)} + 2c s^{-1/3} \mathcal{H}_{(2)} \wedge d\xi], \\ e^{\hat{\phi}} = & \left( \frac{3m}{g^2} \right)^{1/4} s^{-5/6} \Delta^{1/4} X^{-5/6}, \end{aligned} \quad (\text{E.22})$$

where

$$\begin{aligned}
s &\equiv \sin \xi \ , \quad c \equiv \cos \xi \ , \quad X = e^{-\frac{1}{2\sqrt{2}}\phi} \ , \\
\Delta &\equiv 2g X c^2 + 3m X^{-3} s^2 \ , \quad h^i \equiv \sigma^1 - g A_{(1)}^i \ , \\
U &\equiv 9m^2 s^2 X^{-6} - 3g^2 c^2 X^2 + 12mg X^{-2} c^2 - 18mg X^{-2} \ . \quad (\text{E.23})
\end{aligned}$$

## APPENDIX F

FIVE-DIMENSIONAL  $\mathcal{N} = 1$ ,  $SU(2) \times U(1)$ -GAUGED SUPERGRAVITY

The bosonic sector of the  $\mathcal{N} = 4$  gauged supergravity in  $D = 5$  [126] consists of the metric, a scalar, the  $SU(2)$  Yang-Mills potentials  $A_\mu^i$ , a  $U(1)$  gauge potential  $\mathcal{A}_\mu$ , and two 2-form potentials  $B_{\mu\nu}^\alpha$  which transform as a charged doublet under the  $U(1)$ . The Lagrangian is given by

$$\begin{aligned} e^{-1}\mathcal{L} = & -\frac{1}{4}R + \frac{1}{2}(\partial_\mu\phi)(\partial^\mu\phi) - \frac{1}{4}e^{-\frac{8}{\sqrt{6}}\phi}\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu} - \frac{1}{4}e^{\frac{4}{\sqrt{6}}\phi}(F_{\mu\nu}^IF^{I\mu\nu} + B_{\mu\nu}^\alpha B^{\alpha\mu\nu}) \\ & + \frac{1}{8}g_2(g_2e^{-\frac{4}{\sqrt{6}}\phi} + 2\sqrt{2}g_1e^{\frac{2}{\sqrt{6}}\phi}) \\ & + \frac{1}{4}e^{-1}\varepsilon^{\mu\nu\rho\sigma\tau}\left(\frac{1}{g_1}\varepsilon_{\alpha\beta}B_{\mu\nu}^\alpha\mathcal{D}_\rho B_{\sigma\tau}^\beta - F_{\mu\nu}^IF_{\rho\sigma}^I\mathcal{A}^\tau\right), \end{aligned} \quad (\text{F.1})$$

where  $e$  is the determinant of the vielbein,  $g_1$  and  $g_2$  are the gauged couplings of  $U(1)$  and  $SU(2)$  respectively, and  $\varepsilon_{\mu\nu\rho\sigma\tau}$  is a Levi-Civita tensor density. The Abelian field strength  $\mathcal{F}_{\mu\nu}$  and non-Abelian field strength  $F_{\mu\nu}^I$  are given by

$$\mathcal{F}_{\mu\nu} \equiv \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu, \quad F_{\mu\nu}^I \equiv \partial_\mu A_\nu^I - \partial_\nu A_\mu^I + g\varepsilon^{IJK}A_\mu^JA_\nu^K. \quad (\text{F.2})$$

The supersymmetry transformation for the gravitinos is

$$\delta\psi_{\mu a} = \mathcal{D}_\mu\epsilon_a + \Gamma_\mu T_{ab}\epsilon^b - \frac{1}{6\sqrt{2}}(\Gamma_\mu{}^{\nu\rho} - 4\delta_\mu{}^\nu\Gamma^\rho)\left(H_{\nu\rho ab} + \frac{1}{\sqrt{2}}h_{\nu\rho ab}\right)\epsilon^b, \quad (\text{F.3})$$

and that for the gauginos is

$$\delta\chi_a = \frac{1}{\sqrt{2}}\Gamma^\mu\partial_\mu\phi\epsilon_a + A_{ab}\epsilon^b - \frac{1}{2\sqrt{6}}\Gamma^{\mu\nu}(H_{\mu\nu ab} - \sqrt{2}h_{\mu\nu ab})\epsilon^b, \quad (\text{F.4})$$

where  $T_{ab}$ ,  $A_{ab}$ ,  $H_{\mu\nu ab}$  and  $h_{\mu\nu ab}$  are defined as follows

$$T^{ab} \equiv \left(\frac{1}{6\sqrt{2}}g_2e^{-\frac{2}{\sqrt{6}}\phi} + \frac{1}{12}g_1e^{\frac{4}{\sqrt{6}}\phi}\right)(\Gamma_{45})^{ab},$$

$$\begin{aligned}
A^{ab} &\equiv \left( \frac{1}{2\sqrt{6}} g_2 e^{-\frac{2}{\sqrt{6}}\phi} - \frac{1}{2\sqrt{3}} g_1 e^{\frac{4}{\sqrt{6}}\phi} \right) (\Gamma_{45})^{ab} , \\
H_{\mu\nu}^{ab} &\equiv e^{\frac{2}{\sqrt{6}}\phi} [F_{\mu\nu}^I (\Gamma_I)^{ab} + B_{\mu\nu}^\alpha (\Gamma_\alpha)^{ab}] , \quad h_{\mu\nu}^{ab} \equiv e^{-\frac{4}{\sqrt{6}}\phi} \Omega^{ab} \mathcal{F}_{\mu\nu} , 
\end{aligned} \tag{F.5}$$

and  $\Omega^{ab}$  is a charge conjugation matrix. The gauge covariant derivative  $\mathcal{D}_\mu$  acting on Killing spinor is defined by

$$\mathcal{D}_\mu \epsilon_a = \nabla_\mu \epsilon_a + \frac{1}{2} g_1 \mathcal{A}_\mu (\Gamma_{45})_a{}^b \epsilon_b + \frac{1}{2} g_2 A_\mu^I (\Gamma_{45})_a{}^b \epsilon_b , \tag{F.6}$$

with  $\nabla_\mu$  is the usual space-time derivative

$$\nabla_\mu \epsilon_a \equiv \left( \partial_\mu + \frac{1}{4} \omega_\mu^{\alpha\beta} \Gamma_{\alpha\beta} \right) \epsilon_a ,$$

where  $\omega_\mu^{\alpha\beta}$  is the spin connection. Indices  $\alpha$  and  $\beta$  are tangent indices while  $\mu$  and  $\nu$  are space-time indices. The  $\Gamma_{\alpha\beta\dots}$  are the five-dimensional Dirac matrices,

$$\Gamma_{\alpha_1\dots\alpha_n} \equiv \frac{1}{n!} \Gamma_{[\alpha_1} \dots \Gamma_{\alpha_n]} , \quad n = 1, \dots, 5 . \tag{F.7}$$

The Einstein equation of the Lagrangian (F.1) is

$$\begin{aligned}
R_{\mu\nu} &= 2 \partial_\mu \phi \partial_\nu \phi + \frac{1}{6} g_{\mu\nu} (g_2^2 e^{-\frac{4}{\sqrt{6}}\phi} + 2\sqrt{2} g_1 g_2 e^{\frac{2}{\sqrt{6}}\phi}) \\
&\quad + e^{\frac{4}{\sqrt{6}}\phi} \left[ 2 F_{\mu\rho}^I F_\nu{}^\rho + 2 B_{\mu\rho}^\alpha B_\nu{}^\rho - \frac{1}{3} g_{\mu\nu} (F_{\rho\sigma}^I F^{I\rho\sigma} + B_{\rho\sigma}^\alpha B^{\alpha\rho\sigma}) \right] \\
&\quad + e^{-\frac{8}{\sqrt{6}}\phi} \left( 2 \mathcal{F}_{\mu\rho} \mathcal{F}_\nu{}^\rho - \frac{1}{3} g_{\mu\nu} \mathcal{F}_{\rho\sigma} \mathcal{F}^{\rho\sigma} \right) , 
\end{aligned} \tag{F.8}$$

the dilaton equation is

$$\begin{aligned}
\Box \phi &= \left( \frac{1}{2\sqrt{3}} g_1 g_2 e^{\frac{2}{\sqrt{6}}\phi} - \frac{1}{2\sqrt{6}} g_2^2 e^{-\frac{4}{\sqrt{6}}\phi} \right) + \frac{2}{\sqrt{6}} e^{-\frac{8}{\sqrt{6}}\phi} \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} \\
&\quad - \frac{1}{\sqrt{6}} e^{\frac{4}{\sqrt{6}}\phi} (F_{\mu\nu}^I F^{I\mu\nu} + B_{\mu\nu}^\alpha B^{\alpha\mu\nu}) , 
\end{aligned} \tag{F.9}$$

the equations for the gauge fields are

$$\begin{aligned}
\mathcal{D}_\nu(e^{-\frac{8}{\sqrt{6}}\phi} \mathcal{F}^{\nu\mu}) &= \frac{1}{4} e^{-1} \epsilon^{\mu\nu\rho\sigma\tau} (F_{\nu\rho}^I F_{\sigma\tau}^I + B_{\nu\rho}^\alpha B_{\sigma\tau}^\alpha) , \\
\mathcal{D}_\nu(e^{\frac{4}{\sqrt{6}}\phi} F^I{}_{\nu\mu}) &= \frac{1}{2} e^{-1} \epsilon^{\mu\nu\rho\sigma\tau} F_{\nu\rho}^I \mathcal{F}_{\sigma\tau} , \\
e^{\frac{4}{\sqrt{6}}\phi} (B^\alpha)^{\mu\nu} &= \frac{1}{g_1} e^{-1} \epsilon^{\mu\nu\rho\sigma\tau} \epsilon^{\alpha\beta} \mathcal{D}_\rho B_{\sigma\tau}^\beta .
\end{aligned} \tag{F.10}$$

Note that the fields  $B_{(2)}^\alpha$  satisfy a first-order equation of motion, of the kind referred to as “odd-dimensional self-duality” in [137].

As discussed in [126], there are three in-equivalent theories, corresponding to different choices for the gauge couplings in (F.1):  $\mathcal{N} = 4^0$  in which  $g_2 = 0$ ;  $\mathcal{N} = 4^+$  in which  $g_2 = \sqrt{2}g_1$  and  $\mathcal{N} = 4^-$  in which  $g_2 = -\sqrt{2}g_1$  (see also [80]). The  $\mathcal{N} = 4^+$  theory is obtained by truncating the gauged  $SO(6)$ ,  $\mathcal{N} = 8$  supergravity theory in five dimensions, while the  $\mathcal{N} = 4^0$  and  $\mathcal{N} = 4^-$  theories arise as truncations of non-compact  $\mathcal{N} = 8$  supergravities. From the Lagrangian of the Romans’ theory one might conclude that it is not possible to set the  $U(1)$  coupling constant  $g_1$  to be zero. However, as was pointed out in [71], after appropriate rescalings the limit can be taken.

In the language of differential form, the Lagrangian (F.1) can be written as

$$\begin{aligned}
\mathcal{L} = & R * \mathbb{1} - 3X^{-2} * dX \wedge dX - \frac{1}{2} X^4 * \mathcal{F}_{(2)} \wedge \mathcal{F}_{(2)} - \frac{1}{2} X^{-2} (*F_{(2)}^i \wedge F_{(2)}^i + *B_{(2)}^\alpha \wedge B_{(2)}^\alpha) \\
& + \frac{1}{2g_1} \epsilon_{\alpha\beta} B_{(2)}^\alpha \wedge dB_{(2)}^\beta - \frac{1}{2} B_{(2)}^\alpha \wedge B_{(2)}^\alpha \wedge \mathcal{A}_{(1)} - \frac{1}{2} F_{(2)}^i \wedge F_{(2)}^i \wedge \mathcal{A}_{(1)} \\
& + 2g_2(g_2 X^2 + 2\sqrt{2}g_1 X^{-1}) * \mathbb{1} ,
\end{aligned} \tag{F.11}$$

where  $X$  parameterizes the scalar degree of freedom, and can be written in terms of a canonically-normalized dilaton  $\phi$  as  $X = e^{-\frac{1}{\sqrt{6}}\phi}$ . The 2-form field strengths are given by

$$F_{(2)}^i = dA_{(1)}^i + \frac{1}{2} g_2 \epsilon^{ijk} A_{(1)}^j \wedge A_{(1)}^k , \quad \mathcal{F}_{(2)} = d\mathcal{A}_{(1)} . \tag{F.12}$$

Without loss of generality we may set  $g_1 = g$  and  $g_2 = \sqrt{2}g$ , since the two independent gauge coupling constants may be recovered by appropriate rescalings. We also find it advantageous to adopt a complex notation for the two 2-form potentials, which form a charged doublet with respect to the gauge field  $\mathcal{A}_{(1)}$ . Thus we define

$$B_{(2)} \equiv B_{(2)}^1 + i B_{(2)}^2 . \quad (\text{F.13})$$

The equations of motion in differential form notations are

$$\begin{aligned} d(X^{-1} * dX) &= \frac{1}{3} X^4 * \mathcal{F}_{(2)} \wedge \mathcal{F}_{(2)} - \frac{1}{6} X^{-2} (*F_{(2)}^i \wedge F_{(2)}^i + *\bar{B}_{(2)} \wedge B_{(2)}) \\ &\quad - \frac{4}{3} g^2 (X^2 - X^{-1}) * \mathbb{1} , \\ d(X^4 * \mathcal{F}_{(2)}) &= -\frac{1}{2} F_{(2)}^i \wedge F_{(2)}^i - \frac{1}{2} \bar{B}_{(2)} \wedge B_{(2)} , \\ d(X^{-2} * F_{(2)}^i) &= \sqrt{2} g \epsilon^{ijk} X^{-2} * F_{(2)}^j \wedge A_{(1)}^k - F_{(2)}^i \wedge \mathcal{F}_{(2)} , \\ X^2 * F_{(3)} &= -i g B_{(2)} , \\ R_{\mu\nu} &= 3X^{-2} \partial_\mu X \partial_\nu X - \frac{4}{3} g^2 (X^2 + 2X^{-1}) g_{\mu\nu} \\ &\quad + \frac{1}{2} X^4 (\mathcal{F}_\mu{}^\rho \mathcal{F}_{\nu\rho} - \frac{1}{6} g_{\mu\nu} \mathcal{F}_{(2)}^2) + \frac{1}{2} X^{-2} (F_\mu{}^\rho F_{\nu\rho}^i - \frac{1}{6} g_{\mu\nu} (F_{(2)}^i)^2) \\ &\quad + \frac{1}{2} X^{-2} (\bar{B}_{(\mu}{}^\rho B_{\nu)\rho} - \frac{1}{6} g_{\mu\nu} |B_{(2)}|^2) , \end{aligned} \quad (\text{F.14})$$

where

$$F_{(3)} = \mathcal{D} B_{(2)} \equiv dB_{(2)} - i g \mathcal{A}_{(1)} \wedge B_{(2)} . \quad (\text{F.15})$$

The operator  $\mathcal{D}$  defined in this equation is the  $U(1)$  gauge-covariant exterior derivative.



## VITA

Tuan Anh Tran received a Bachelor of Science in Physics from Hanoi University, Vietnam, in July 1992. From 1993 to 1994, he was a graduate student at the Institute of Physics, National Center for Science and Technology, Hanoi, Vietnam. He joined the Diploma Program at the Abdus Salam International Center for Theoretical Physics (ICTP), Trieste, Italy in September 1994 and completed the program in August 1995. After spending most of his time from September 1995 to August 1996 as a visiting researcher at ICTP, he joined the graduate program at Texas A&M University in September 1996. His address is the Institute for Fundamental Theory, University of Florida, P.O. Box 118440, Gainesville, FL 32611-8440.

The typist for this thesis was Tuan A. Tran.